

Reference

American Association of Cereal Chemists (AACC). (1999). American Association of Cereal Chemists Approved Methods, 16–18.

American Association of Cereal Chemists (AACC). (2001). The Definition of Dietary Fiber, 112(3).

Retrieved from

<https://www.aaccnet.org/initiatives/definitions/Documents/DietaryFiber/DFDef.pdf>

ANDERSON, N. E., & YDESCALE, F. M. C. (1980). AN ANALYSIS OF THE DIETARY FIBER, CONTENT OF A STANDARD WHEAT BRAN. *Journal of Food Science*, 45(2), 336–340.

<http://doi.org/10.1111/j.1365-2621.1980.tb02609.x>

Badan Pengawas Obat dan Makanan Republik Indonesia (BPOM). (2016a). PERATURAN KEPALA BADAN PENGAWAS OBAT DAN MAKANAN REPUBLIK INDONESIA NOMOR 13 TAHUN 2016 TENTANG PENGAWASAN KLAIM PADA LABEL DAN IKLAN PANGAN OLAHAN.

Badan Pengawas Obat dan Makanan Republik Indonesia (BPOM). (2016b). PerKa BPOM no 21 tahun 2016 Tentang Kategori Pangan, 1–28.

Chen, J. S., Fei, M. J., Shi, C. L., Tian, J. C., Sun, C. L., Zhang, H., ... Dong, H. X. (2011). Effect of particle size and addition level of wheat bran on quality of dry white Chinese noodles. *Journal of Cereal Science*, 53(2), 217–224. <http://doi.org/10.1016/j.jcs.2010.12.005>

Czernhorsky, J. H., & Hooker, R. (2008). The chemistry of baking. *NZ Institute for Crop and Food Research*, 1–8.

Fu, B. X. (2008). Asian noodles: History, classification, raw materials, and processing. *Food Research International*, 41(9), 888–902. <http://doi.org/10.1016/j.foodres.2007.11.007>

Gulia, N., Dhaka, V., & Khatkar, B. S. (2014). Instant Noodles: Processing, Quality, and Nutritional Aspects. *Critical Reviews in Food Science and Nutrition*, 54(10), 1386–1399.
<http://doi.org/10.1080/10408398.2011.638227>

Hatcher, D. W., Anderson, M. J., Desjardins, R. G., Edwards, N. M., & Dexter, J. E. (2002). Effects of flour particle size and starch damage on processing and quality of white salted noodles. *Cereal*

- Chemistry*, 79(1), 64–71. <http://doi.org/10.1094/CCHEM.2002.79.1.64>
- Hemdane, S., Jacobs, P. J., Dornez, E., Verspreet, J., Delcour, J. A., & Courtin, C. M. (2016). Wheat (Triticum aestivum L.) Bran in Bread Making: A Critical Review. *Comprehensive Reviews in Food Science and Food Safety*, 15(1), 28–42. <http://doi.org/10.1111/1541-4337.12176>
- Hipsley, E. (1953). Dietary “Fibre” and pregnancy toxæmia. *Br Med J*, (2), 420–422.
- Izydorczyk, M. S., Lagassé, S. L., Hatcher, D. W., Dexter, J. E., & Rossnagel, B. G. (2005). The enrichment of Asian noodles with fiber-rich fractions derived from roller milling of hull-less barley. *Journal of the Science of Food and Agriculture*, 85(12), 2094–2104.
<http://doi.org/10.1002/jsfa.2242>
- Kubomura, K. (1998). Instant noodles in Japan. *Cereal Foods World*, (43), 194–197.
- Lawless, H. T., & Heyman, H. (2010). *Sensory Evaluation of Food*. (D. R. Heldman, Ed.) (Second).
- Lehtinen, O. (2012). Modifying Wheat Bran for Food Applications - Effect of Wet Milling and Enzymatic Treatment, (May), 91.
- Ma, D. Y., Zhang, J., Lou, Wang, X. N., Wang, C. Y., & Guo, T. C. (2014). Color, cooking properties and texture of yellow alkaline noodles enriched with millet and corn flour. *International Food Research Journal*, 21(3), 1187–1192. Retrieved from <http://www.ifrj.upm.edu.my>
- Maes, C., & Delcour, J. A. (2002). Structural characterisation of water extractable and water unextractable arabinoxylans in wheat bran. *Journal of Cereal Science*, 35, 315–326.
- Menteri Kesehatan Republik Indonesia. (2014). PERATURAN MENTERI KESEHATAN REPUBLIK INDONESIA NOMOR 41 TAHUN 2014 TENTANG PEDOMAN GIZI SEIMBANG. Retrieved from http://www.hukor.depkes.go.id/uploads/produk_hukum/PMK No. 41 ttg Pedoman Gizi Seimbang.pdf
- Niu, M., Hou, G. G., Kindelspire, J., Krishnan, P., & Zhao, S. (2017). Microstructural, textural, and sensory properties of whole-wheat noodle modified by enzymes and emulsifiers. *Food Chemistry*, 223, 16–24. <http://doi.org/10.1016/j.foodchem.2016.12.021>
- Niu, M., Hou, G. G., Wang, L., & Chen, Z. (2014). Effects of superfine grinding on the quality

- characteristics of whole-wheat flour and its raw noodle product. *Journal of Cereal Science*, 60(2), 382–388. <http://doi.org/10.1016/j.jcs.2014.05.007>
- Pakhare, K., Dagadkhair, A., Udachan, I., & Andhale, R. (2016). Studies on Preparation and Quality of Nutritious Noodles by Incorporation of Defatted Rice Bran and Soy Flour. *J Food Process Technol*, 7(10). <http://doi.org/10.4172/2157-7110.1000629>
- Prairahong, P. (2002). *FORMULATION OF DIETARY FIBER-ENRICHED INSTANT NOODLES*. Chulalongkorn University.
- Reungmaneepaitoon, S., & Sikkhamondhol, C. (2014). Nutritive improvement of instant fried noodles with oat bran Nutritive improvement of instant fried noodles with oat bran, (March 2006).
- Reungmaneepaitoon, S., Sikkhamondhol, C., & Tiangpook, C. (2006). Nutritive improvement of instant fried noodles with oat bran. *Songklanakarin Journal of Science and Technology*, 28(SUPPL. 1), 89–97.
- Ritthiruangdej, P., Parnbankled, S., Donchedee, S., & Wongsagonsup, R. (2011). Physical, Chemical, Textural and Sensory Properties of Dried Wheat Noodles Supplemented with Unripe Banana Flour. *Kasetsart J. (Nat. Sci.)*, 45, 500–509. Retrieved from <https://pdfs.semanticscholar.org/b7d0/ddd1f0a9c0182068a011c66307f9cad4ed05.pdf>
- Song, X., Zhu, W., Pei, Y., Ai, Z., & Chen, J. (2013). Effects of wheat bran with different colors on the qualities of dry noodles. *Journal of Cereal Science*, 58(3), 400–407.
<http://doi.org/10.1016/j.jcs.2013.08.005>
- Stevenson, L., & et.al. (2012). Wheat bran: its composition and benefits to health, a European perspective. *International Journal of Food Sciences and Nutrition*, 63(8), 1001–1013.
<http://doi.org/10.3109/09637486.2012.687366>
- Stewart, N. (2014). The health benefits of dietary fiber consumption of adults in the United States.
- Tan, T. C., Phatthanawiboon, T., & Mat Easa, A. (2016). Quality, Textural, and Sensory Properties of Yellow Alkaline Noodles Formulated with Salted Duck Egg White. *Journal of Food Quality*, 39(4), 342–350. <http://doi.org/10.1111/jfq.12203>

- Trowell, H., Burkitt, D., & Heaton, K. (1985). Definitions of dietary fibre and fibre-depleted foods and disease., 21–30.
- USAID. (1970). Micronutrient Content of Wheat and Wheat Flour, 1–4. Retrieved from
<http://linkinghub.elsevier.com/retrieve/pii/S018972411530076X>
- Vernaza, M. G., Gualarte, M. A., & Chang, Y. K. (2011). Addition of Green Banana Flour to Instant Noodles: Rheological and technological properties. *Ciencia e Agrotecnologia*, 35(6), 1157–1165.
<http://doi.org/10.1590/S1413-70542011000600016>
- Wheat Marketing Center. (2008). *Wheat and flour testing methods : A guide to understanding wheat and flour quality* (2nd ed.). Kansas State University.
- WHO. (2017). WHO | NCD mortality and morbidity. WHO. Retrieved from
http://www.who.int/gho/ncd/mortality_morbidity/en/
- Wichchukit, S., & O'Mahony, M. (2015). The 9-point hedonic scale and hedonic ranking in food science: Some reappraisals and alternatives. *Journal of the Science of Food and Agriculture*, 95(11), 2167–2178. <http://doi.org/10.1002/jsfa.6993>
- Yun, S.-H., Rema, G., & Quail, K. (1997). Instrumental assessments of Japanese white salted noodle quality. *J. Sci. Food Agric*, (74), 81–88.
- Zhang, N., & Ma, G. (2016). Noodles, traditionally and today. *Journal of Ethnic Foods*, 3(3), 209–212.
<http://doi.org/10.1016/j.jef.2016.08.003>