

## REFERENCES

- Amirudin, A., Urbański, M., Saputra, J., Johansyah, M., Latip, L., Tarmizi, A., & Afrizal, T. (2021). The Impact of the COVID-19 Self-Isolation Policy on the Occupations of Vulnerable Groups. *International Journal Of Environmental Research And Public Health*, 18(12), 6452. doi: 10.3390/ijerph18126452
- Armbruster, D. A., & Pry, T. (2008). Limit of blank, limit of detection and limit of quantitation. *The Clinical biochemist. Reviews*, 29 Suppl 1(Suppl 1), S49–S52.
- Batéjat, C., Grassin, Q., Manuguerra, J., & Leclercq, I. (2021). Heat inactivation of the severe acute respiratory syndrome coronavirus 2. *Journal Of Biosafety And Biosecurity*, 3(1), 1-3. <https://doi.org/10.1016/j.jobb.2020.12.001>
- Ben-Assa, N., Naddaf, R., Gefen, T., Capucha, T., Hajjo, H., & Mandelbaum, N. et al. (2020). Direct on-the-spot detection of SARS-CoV-2 in patients. *Experimental Biology And Medicine*, 245(14), 1187-1193. <https://doi.org/10.1177/1535370220941819>
- Beyene, G., Alemu, F., Kebede, E., Alemayehu, D., Seyoum, T., & Tefera, D. et al. (2021). Saliva is superior over nasopharyngeal swab for detecting SARS-CoV2 in COVID-19 patients. *Scientific Reports*, 11(1). <https://doi.org/10.1038/s41598-021-02097-2>
- Biolabs, N. (2022). International.neb.com. Retrieved 1 June 2022, from <https://international.neb.com/products/m1800-warmstart-colorimetric-lamp-2x-master-mix-dna-rna#Product%20Information>.
- Burns, J., Butler, J., Moran, J., & Whitesides, G. (1991). Selective reduction of disulfides by tris(2-carboxyethyl)phosphine. *The Journal Of Organic Chemistry*, 56(8), 2648-2650. <https://doi.org/10.1021/jo00008a014>
- Bustin, S., & Nolan, T. (2020). RT-qPCR Testing of SARS-CoV-2: A Primer. *International Journal Of Molecular Sciences*, 21(8), 3004. <https://doi.org/10.3390/ijms21083004>
- Chen, J., Yip, C., Poon, R., Chan, K., Cheng, V., & Hung, I. et al. (2020). Evaluating the use of posterior oropharyngeal saliva in a point-of-care assay for the detection of SARS-CoV-2. *Emerging Microbes & Infections*, 9(1), 1356-1359. <https://doi.org/10.1080/22221751.2020.1775133>
- Cook, B., Kobasa, K., Tamayo, M., Theriault, N., Gordon Pappas, D., & Theriault, S. (2022). Implementation of a Rapid RT-LAMP Saliva-Based SARS-CoV-2 Testing Program in the Workplace. *Diagnostics*, 12(2), 474. <https://doi.org/10.3390/diagnostics12020474>
- Dao Thi, V., Herbst, K., Boerner, K., Meurer, M., Kremer, L., & Kirrmaier, D. et al. (2020). A colorimetric RT-LAMP assay and LAMP-sequencing for detecting SARS-CoV-2 RNA in clinical samples. *Science Translational Medicine*, 12(556), eabc7075. <https://doi.org/10.1126/scitranslmed.abc7075>
- Dhar Chowdhury, S., & Oommen, A. (2020). Epidemiology of COVID-19. *Journal Of Digestive Endoscopy*, 11(01), 03-07. doi: 10.1055/s-0040-1712187
- Farrell, R. (2017). Creating a Ribonuclease-Free Environment. *RNA Methodologies*, 55-74.

<https://doi.org/10.1016/b978-0-12-804678-4.00002-6>

- Ganguli, A., Mostafa, A., Berger, J., Lim, J., Araud, E., & Baek, J. et al. (2021). Reverse Transcription Loop-Mediated Isothermal Amplification Assay for Ultrasensitive Detection of SARS-CoV-2 in Saliva and Viral Transport Medium Clinical Samples. *Analytical Chemistry*, 93(22), 7797-7807. doi: 10.1021/acs.analchem.0c05170
- Gloster, A., Lamnisos, D., Lubenko, J., Presti, G., Squatrito, V., & Constantinou, M. et al. (2020). Impact of COVID-19 pandemic on mental health: An international study. *PLOS ONE*, 15(12), e0244809. doi: 10.1371/journal.pone.0244809
- Hallmaier-Wacker, L., Lueert, S., Roos, C., & Knauf, S. (2018). The impact of storage buffer, DNA extraction method, and polymerase on microbial analysis. *Scientific Reports*, 8(1). <https://doi.org/10.1038/s41598-018-24573-y>
- Hayden, A., Kuentzel, M., & Chittur, S. (2021). Rapid, Affordable, and Scalable SARS-CoV-2 Detection From Saliva. *Journal Of Biomolecular Techniques : JBT*, 32(3), 148-157. doi: 10.7171/jbt.21-3203-010
- Hölz, K., Pavlic, A., Lietard, J., & Somoza, M. (2019). Specificity and Efficiency of the Uracil DNA Glycosylase-Mediated Strand Cleavage Surveyed on Large Sequence Libraries. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-54044-x>
- Huang, X., Tang, G., Ismail, N., & Wang, X. (2022). Developing RT-LAMP assays for rapid diagnosis of SARS-CoV-2 in saliva. *Ebiomedicine*, 75, 103736. doi: 10.1016/j.ebiom.2021.103736
- Kobayashi, G., Brito, L., Moreira, D., Suzuki, A., Hsia, G., & Pimentel, L. et al. (2021). A Novel Saliva RT-LAMP Workflow for Rapid Identification of COVID-19 Cases and Restraining Viral Spread. *Diagnostics*, 11(8), 1400. <https://doi.org/10.3390/diagnostics11081400>
- Koczula, K., & Gallotta, A. (2016). Lateral flow assays. *Essays In Biochemistry*, 60(1), 111-120. doi: 10.1042/ebc20150012
- Kralik, P., & Ricchi, M. (2017). A Basic Guide to Real Time PCR in Microbial Diagnostics: Definitions, Parameters, and Everything. *Frontiers In Microbiology*, 8. doi: 10.3389/fmicb.2017.00108
- Kundrod, K., Natoli, M., Chang, M., Smith, C., Paul, S., & Ogoe, D. et al. (2022). Sample-to-answer, extraction-free, real-time RT-LAMP test for SARS-CoV-2 in nasopharyngeal, nasal, and saliva samples: Implications and use for surveillance testing. *PLOS ONE*, 17(2), e0264130. <https://doi.org/10.1371/journal.pone.0264130>
- Lamb, L., Bartolone, S., Ward, E., & Chancellor, M. (2020). Rapid detection of novel coronavirus/Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) by reverse transcription-loop-mediated isothermal amplification. *PLOS ONE*, 15(6), e0234682. <https://doi.org/10.1371/journal.pone.0234682>
- Lalli, M., Langmade, J., Chen, X., Fronick, C., Sawyer, C., & Burcea, L. et al. (2020). Rapid and Extraction-Free Detection of SARS-CoV-2 from Saliva by Colorimetric Reverse-Transcription Loop-Mediated Isothermal Amplification. *Clinical Chemistry*, 67(2), 415-424. doi: 10.1093/clinchem/hvaa267

- Liu, X., Feng, J., Zhang, Q., Guo, D., Zhang, L., & Suo, T. et al. (2020). Analytical comparisons of SARS-CoV-2 detection by qRT-PCR and ddPCR with multiple primer/probe sets. *Emerging Microbes & Infections*, 9(1), 1175-1179. <https://doi.org/10.1080/22221751.2020.1772679>
- Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., & Wu, H. et al. (2020). Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *The Lancet*, 395(10224), 565-574. doi: 10.1016/s0140-6736(20)30251-8
- Maxim, L., Niebo, R., & Utell, M. (2014). Screening tests: a review with examples. *Inhalation Toxicology*, 26(13), 811-828. doi: 10.3109/08958378.2014.955932
- Mendelson, Y. (2012). Biomedical Sensors. *Introduction To Biomedical Engineering*, 609-666. <https://doi.org/10.1016/b978-0-12-374979-6.00010-1>
- Mohamadian, M., Chiti, H., Shoghli, A., Biglari, S., Parsamanesh, N., & Esmailzadeh, A. (2021). COVID-19: Virology, biology and novel laboratory diagnosis. *The Journal Of Gene Medicine*, 23(2). doi: 10.1002/jgm.3303
- Mohd Thabit, A., Peariasamy, K., Kuan, P., Fern Ying, D., Nheu, N., & Cyncynatus, C. et al. (2021). Diagnostic accuracy of fresh drooled saliva for SARS-CoV-2 in travelers. *Travel Medicine And Infectious Disease*, 43, 102144. <https://doi.org/10.1016/j.tmaid.2021.102144>
- Newman, C., Ramuta, M., McLaughlin, M., Wiseman, R., Karl, J., & Dudley, D. et al. (2021). Initial Evaluation of a Mobile SARS-CoV-2 RT-LAMP Testing Strategy. *Journal Of Biomolecular Techniques : JBT*, 32(3), 137-147. <https://doi.org/10.7171/jbt.21-32-03-009>
- Notomi, T., Okayama, H., Masubuchi, H., Yonekawa, T., Watanabe, K., Amino, N., & Hase, T. (2000). Loop-mediated isothermal amplification of DNA. *Nucleic acids research*, 28(12), E63. <https://doi.org/10.1093/nar/28.12.e63>
- Panda, B., Meher, A., & Hazra, R. (2019). Comparison between different methods of DNA isolation from dried blood spots for determination of malaria to determine specificity and cost effectiveness. *Journal Of Parasitic Diseases*, 43(3), 337-342. <https://doi.org/10.1007/s12639-019-01136-0>
- Qamar, W., Khan, M., & Arafah, A. (2017). Optimization of conditions to extract high quality DNA for PCR analysis from whole blood using SDS-proteinase K method. *Saudi Journal Of Biological Sciences*, 24(7), 1465-1469. <https://doi.org/10.1016/j.sjbs.2016.09.016>
- Reolo, M., Eleazar, C., Sonio, J., Solon, R., Villamor, J., Loedin, A., & Moore, K. (2021). Saliva “Treat-and-Heat” Triplex Reverse Transcription Loop-Mediated Isothermal Amplification Assay for SARS-CoV-2. *Journal Of Biomolecular Techniques : JBT*, 32(3), 186-198. doi: 10.7171/jbt.21-3203-014
- Sahoo, P., Sethy, K., Mohapatra, S., & Panda, D. (2016). Loop mediated isothermal amplification: An innovative gene amplification technique for animal diseases. *Veterinary World*, 9(5), 465-469. doi: 10.14202/vetworld.2016.465-469
- Tanner, N., Zhang, Y., & Evans, T. (2012). Simultaneous multiple target detection in real-time loop-mediated isothermal amplification. *Biotechniques*, 53(2), 81-89. <https://doi.org/10.2144/0000113902>

- Udugama, B., Kadhiresan, P., Kozlowski, H., Malekjahani, A., Osborne, M., & Li, V. et al. (2020). Diagnosing COVID-19: The Disease and Tools for Detection. *ACS Nano*, *14*(4), 3822-3835. doi: 10.1021/acsnano.0c02624
- Urrutia-Cabrera, D., Liou, R., Wang, J., Chan, J., Hung, S., & Hewitt, A. et al. (2021). Comparative analysis of loop-mediated isothermal amplification (LAMP)-based assays for rapid detection of SARS-CoV-2 genes. *Scientific Reports*, *11*(1). <https://doi.org/10.1038/s41598-021-01472-3>
- Vandenberg, O., Martiny, D., Rochas, O., van Belkum, A., & Kozlakidis, Z. (2020). Considerations for diagnostic COVID-19 tests. *Nature Reviews Microbiology*, *19*(3), 171-183. doi: 10.1038/s41579-020-00461-z
- Vaz, S., Santana, D., Netto, E., Pedroso, C., Wang, W., Santos, F., & Brites, C. (2020). Saliva is a reliable, non-invasive specimen for SARS-CoV-2 detection. *The Brazilian Journal Of Infectious Diseases*, *24*(5), 422-427. <https://doi.org/10.1016/j.bjid.2020.08.001>
- Wei, S., Suryawanshi, H., Djandji, A., Kohl, E., Morgan, S., & Hod, E. et al. (2021). Field-deployable, rapid diagnostic testing of saliva for SARS-CoV-2. *Scientific Reports*, *11*(1). doi: 10.1038/s41598-021-84792-8
- Wu, F., Zhao, S., Yu, B., Chen, Y., Wang, W., & Song, Z. et al. (2020). A new coronavirus associated with human respiratory disease in China. *Nature*, *579*(7798), 265-269. <https://doi.org/10.1038/s41586-020-2008-3>
- Yamazaki, W., Matsumura, Y., Thongchankaew-Seo, U., Yamazaki, Y., & Nagao, M. (2021). Development of a point-of-care test to detect SARS-CoV-2 from saliva which combines a simple RNA extraction method with colorimetric reverse transcription loop-mediated isothermal amplification detection. *Journal Of Clinical Virology*, *136*, 104760. doi: 10.1016/j.jcv.2021.104760
- Yang, Q., Meyerson, N., Clark, S., Paige, C., Fattor, W., & Gilchrist, A. et al. (2021). Saliva TwoStep for rapid detection of asymptomatic SARS-CoV-2 carriers. *Elife*, *10*. <https://doi.org/10.7554/elife.65113>
- Yu, A., Galatsis, K., Zheng, J., Le, J., Ma, D., Perlman, S., & Rosbash, M. (2021). Development of a Saliva-Optimized RT-LAMP Assay for SARS-CoV-2. *Journal Of Biomolecular Techniques : JBT*, *32*(3), 102-113. doi: 10.7171/jbt.21-3203-005
- Yu, F., Yan, L., Wang, N., Yang, S., Wang, L., & Tang, Y. et al. (2020). Quantitative Detection and Viral Load Analysis of SARS-CoV-2 in Infected Patients. *Clinical Infectious Diseases*, *71*(15), 793-798. doi: 10.1093/cid/cia345
- Zhang, Q., McGuigan, C., Lew, K., & Chris Le, X. (2012). Urine Sample Collection and Handling. *Comprehensive Sampling And Sample Preparation*, 123-142. <https://doi.org/10.1016/b978-0-12-381373-2.00069-7>
- Zhang, X., Huang, H., Zhuang, D., Nasser, M., Yang, M., Zhu, P., & Zhao, M. (2020). Biological, clinical and epidemiological features of COVID-19, SARS and MERS and AutoDock simulation of ACE2. *Infectious Diseases Of Poverty*, *9*(1). doi: 10.1186/s40249-020-00691-6