

## REFERENCES

- A, D. U., E, O. S., & A, O. N. (2020). Antibiogram of bacteria isolated from roasted bush meat and chicken samples sold in Afikpo, Nigeria. International Journal of Psychosocial Rehabilitation, 24(04), 848–861. <https://doi.org/10.37200/ijpr/v24i4/pr201058>
- Alatorre-Meda, M., Taboada, P., Hartl, F., Wagner, T., Freis, M., & Rodríguez, J. R. (2011). The influence of chitosan valence on the complexation and transfection of DNA: The weaker the DNA–chitosan binding the higher the transfection efficiency. Colloids and Surfaces B: Biointerfaces, 82(1), 54–62. <https://doi.org/10.1016/j.colsurfb.2010.08.013>
- Aliño, M., Grau, R., Toldrá, F., Blesa, E., Pagán, M. J., & Barat, J. M. (2009). Influence of sodium replacement on physicochemical properties of dry-cured loin. Meat Science, 83(3), 423–430. <https://doi.org/10.1016/j.meatsci.2009.06.022>
- Almashhadany, D. A. (2019). Occurrence and antimicrobial susceptibility of salmonella isolates from grilled chicken meat sold at retail outlets in Erbil City, Kurdistan Region, Iraq. Italian Journal of Food Safety, 8(2). <https://doi.org/10.4081/ijfs.2019.8233>
- Arabyan, N., Huang, B.C. and Weimer, B.C. (2017) ‘Amylases and their importance during glycan degradation: Genome sequence release of&nbs; &nbs; &nbs; &nbs; &nbs; &nbs; salmonella&nbs; &nbs; &nbs; &nbs; &nbs; amylase knockout strains’, Genome Announcements, 5(20). doi:10.1128/genomea.00355-17.
- Batista, A. C. L., Dantas, G. C., Santos, J., & Amorim, R. V. S. (2011). Antimicrobial effects of native chitosan against opportunistic gram-negative bacteria. Microbiology Journal, 1(3), 105–112. <https://doi.org/10.3923/mj.2011.105.112>
- Boeriu, C. G., & Broek, L. A. M. (2019). Chemical and enzymatic modification of chitosan to produce new functional materials with improved properties. Chitin and Chitosan, 245–258. <https://doi.org/10.1002/9781119450467.ch10>
- Chignola, R., Mainente, F., & Zoccatelli, G. (2022). Rheology of individual chitosan and polyphenol/Chitosan microparticles for Food Engineering. Food Hydrocolloids, 132, 107869. <https://doi.org/10.1016/j.foodhyd.2022.107869>
- Chumngoen, W., Chen, H.-Y., & Tan, F.-J. (2016). Validation of feasibility and quality of chicken breast meat cooked under various water-cooking conditions. Animal Science Journal, 87(12), 1536–1544. <https://doi.org/10.1111/asj.12589>

- Comaposada, J., Arnau, J., & Gou, P. (2007). Sorption isotherms of salted minced pork and of lean surface of dry-cured hams at the end of the resting period using KCl as substitute for NaCl. Meat Science, 77(4), 643–648. <https://doi.org/10.1016/j.meatsci.2007.05.017>
- Creasy, A., Barker, G., & Carta, G. (2017). Systematic interpolation method predicts protein chromatographic elution with salt gradients, pH gradients and combined salt/pH gradients. Biotechnology Journal, 12(3), 1600636. <https://doi.org/10.1002/biot.201600636>
- Darmadji, P., & Izumimoto, M. (1994). Effect of chitosan in meat preservation. Meat Science, 38(2), 243–254. [https://doi.org/10.1016/0309-1740\(94\)90114-7](https://doi.org/10.1016/0309-1740(94)90114-7)
- Dessy Setiawanti. (2021). Implementasi 2d Barcode Pada Label Pangan Olahan Sebagai kekuatan pengawasan Badan Pengawas obat Dan Makanan (BPOM). Food Scientia : Journal of Food Science and Technology, 1(2), 87–102. <https://doi.org/10.33830/fsj.v1i2.1968.2021>
- Ding, X., Wang, G., Zou, Y., Zhao, Y., Ge, C., & Liao, G. (2021). Evaluation of small molecular metabolites and sensory properties of Xuanwei Ham salted with partial replacement of NaCl by KCl. Meat Science, 175, 108465. <https://doi.org/10.1016/j.meatsci.2021.108465>
- FDA. (2020). 2018 NARMS update: Integrated Report Summary Interactive Version. U.S. Food and Drug Administration. <https://www.fda.gov/animal-veterinary/national-antimicrobial-resistance-monitoring-system/2018-narms-update-integrated-report-summary-interactive-version>
- Goo, B.G. and Park, J.K. (2020) ‘Three-dimensional structure analysis of affinity between enzyme and substrate affecting on enzyme activity’, Journal of Chitin and Chitosan, 25(3), pp. 117–125. doi:10.17642/jcc.25.3.2.
- Goli, T., Bohuon, P., Ricci, J., & Colligan, A. (2012). Evolution of pH during immersion of meat protein matrices in acidic marinades. Meat Science, 90(3), 618–623. <https://doi.org/10.1016/j.meatsci.2011.10.003>
- HAND, L. W., TERRELL, R. N., & SMITH, G. C. (1982). Effects of complete or partial replacement of sodium chloride on processing and sensory properties of hams. Journal of Food Science, 47(6), 1776–1778. <https://doi.org/10.1111/j.1365-2621.1982.tb12880.x>
- Jung, Y.-K. et al. (2011) “Comparison of quality traits of thigh meat from Korean native chickens and broilers,” Korean Journal for Food Science of Animal Resources, 31(5), pp. 684–692. Available at: <https://doi.org/10.5851/kosfa.2011.31.5.684>.
- Karam, L., & Mallah, A. (2023). Application of chitosan in active food packaging. Chitosan: Novel Applications in Food Systems, 1–30. <https://doi.org/10.1016/b978-0-12-821663-7.00002-8>

- Kim, H.-W., Hwang, K.-E., Song, D.-H., Kim, Y.-J., Ham, Y.-K., Yeo, E.-J., Jeong, T.-J., Choi, Y.-S., & Kim, C.-J. (2015). Effect of pre-rigor salting levels on physicochemical and textural properties of chicken breast muscles. *Korean Journal for Food Science of Animal Resources*, 35(5), 577–584. <https://doi.org/10.5851/kosfa.2015.35.5.577>
- Kralik, G., Kralik, Z., Grčević, M., & Hanžek, D. (2018). Quality of chicken meat. *Animal Husbandry and Nutrition*. <https://doi.org/10.5772/intechopen.72865>
- Lähteenmäki, K. et al. (2004) ‘Antiprotease inactivation by salmonella enterica released from infected macrophages’, *Cellular Microbiology*, 7(4), pp. 529–538. doi:10.1111/j.1462-5822.2004.00483.x.
- Leygonie, C., Britz, T.J. and Hoffman, L.C. (2012) “Impact of freezing and thawing on the quality of meat: Review,” *Meat Science*, 91(2), pp. 93–98. Available at: <https://doi.org/10.1016/j.meatsci.2012.01.013>.
- Limbardon, P., Lubis, N. D., Amelia, R., & Amelia, S. (2022). Food handling practice of culinary merchants in Medan, Indonesia. *Journal of Asian Multicultural Research for Medical and Health Science Study*, 3(1), 36–46. <https://doi.org/10.47616/jamrmhss.v3i1.201>
- Liu, X.D. et al. (2012) “Differential proteome analysis of breast and thigh muscles between Korean native chickens and commercial broilers,” *Asian-Australasian Journal of Animal Sciences*, 25(6), pp. 895–902. Available at: <https://doi.org/10.5713/ajas.2011.11374>.
- Lopez, K., Schilling, M. W., Armstrong, T. W., Smith, B. S., & Corzo, A. (2012). Sodium chloride concentration affects yield, quality, and sensory acceptability of vacuum-tumbled marinated broiler breast fillets. *Poultry Science*, 91(5), 1186–1194. <https://doi.org/10.3382/ps.2011-01733>
- Manoj, J. (2015). The role of poultry in Food Borne Salmonellosis and its public health importance. *Advances in Animal and Veterinary Sciences*, 3(9), 485–490. <https://doi.org/10.14737/journal.aavs/2015/3.9.485.490>
- Menconi, A., Pumford, N. R., Morgan, M. J., Bielke, L. R., Kallapura, G., Latorre, J. D., Wolfenden, A. D., Hernandez-Velasco, X., Hargis, B. M., & Tellez, G. (2014). Effect of chitosan on salmonella typhimurium in broiler chickens. *Foodborne Pathogens and Disease*, 11(2), 165–169. <https://doi.org/10.1089/fpd.2013.1628>
- Miranda, L.T., Rakovski, C. and Were, L.M. (2012) “Effect of maillard reaction products on oxidation products in ground chicken breast,” *Meat Science*, 90(2), pp. 352–360. Available at: <https://doi.org/10.1016/j.meatsci.2011.07.022>.

Montal, M. (1999) 'Electrostatic attraction at the core of Membrane Fusion', FEBS Letters, 447(2–3), pp. 129–130. doi:10.1016/s0014-5793(99)90269-7.

Mostafa, T.B. and Darwish, A.S. (2014) 'An approach toward construction of tuned chitosan/polyaniline/metal hybrid nanocomposites for treatment of meat industry wastewater', Chemical Engineering Journal, 243, pp. 326–339. doi:10.1016/j.cej.2014.01.006.

Nahar, M. K., Zakaria, Z., & Hashim, U. (2013). Effect of buffer and ph on the protein extraction for chicken meat. Advanced Materials Research, 795, 206–210. <https://doi.org/10.4028/www.scientific.net/amr.795.206>

Norris, V. (2019) 'Faculty opinions recommendation of threonine phosphorylation fine-tunes the regulatory activity of histone-like nucleoid structuring protein in salmonella transcription.', Faculty Opinions – Post-Publication Peer Review of the Biomedical Literature [Preprint]. doi:10.3410/f.736245481.793563767.

Okawa, S. (2012) 'Suppressing method of supercooling state in cool box using membrane', Supercooling [Preprint]. doi:10.5772/35545.

Oscar, T.P. (2014a) 'General regression neural network model for behavior of salmonella on chicken meat during Cold storage', Journal of Food Science, 79(5). doi:10.1111/1750-3841.12435.

PÄRN, T., DAHL, V., LIENEMANN, T., PEREVOSČIKOV, J., & DE JONG, B. (2017). Multi-country outbreak of salmonella enteritidis infection linked to the International Ice Hockey Tournament. Epidemiology and Infection, 145(11), 2221–2230. <https://doi.org/10.1017/s0950268817001212>

Park, S., Marsh, K. S., & Dawson, P. (2010). Application of chitosan-incorporated LDPE film to Sliced Fresh Red Meats for shelf life extension. Meat Science, 85(3), 493–499. <https://doi.org/10.1016/j.meatsci.2010.02.022>

Podolak, R., Enache, E., Stone, W., Black, D. G., & Elliott, P. H. (2010). Sources and risk factors for contamination, survival, persistence, and heat resistance of salmonella in low-moisture foods. Journal of Food Protection, 73(10), 1919–1936. <https://doi.org/10.4315/0362-028x-73.10.1919>

Popa, G. L., & Popa, M. I. (2021). Salmonella spp. infection – a continuous threat worldwide. GERMS, 11(1), 88–96. <https://doi.org/10.18683/germs.2021.1244>

Purnomo, D. et al. (2018) 'The mapping of halal critical points in the process of distribution chain of chicken meat products in West Java (case study: Traditional slaughtering house in Y Chicken House west java)', KnE Life Sciences, 4(2), p. 9. doi:10.18502/cls.v4i2.1652.

Purwantoro, A., Drastini, Y., Khasanah, H., Qadrianti, H., Indiana, E. D., & Artama, W. T. (2019). Micro-business of chicken meat trading at Yogyakarta city traditional markets, Indonesia. Proceedings of the International Conference of Business, Economy, Entrepreneurship and Management. <https://doi.org/10.5220/0009961706590661>

Qiao, M., Fletcher, D. L., Smith, D. P., & Northcutt, J. K. (2001). The effect of broiler breast meat color on pH, moisture, water-holding capacity, and emulsification capacity. *Poultry Science*, 80(5), 676–680. <https://doi.org/10.1093/ps/80.5.676>

Rahman, M. S., & Rahman, M. R. (2020). Ph in food preservation. *Handbook of Food Preservation*, 323–332. <https://doi.org/10.1201/9780429091483-23>

Rapa, M., & Vasile, C. (2020). Chitin- and chitosan-based Bionanocomposites for active packaging. *Chitin- and Chitosan-Based Biocomposites for Food Packaging Applications*, 59–68. <https://doi.org/10.1201/9780429299605-6>

Roy, P., Dhillon, A. S., Lauerman, L. H., Schaberg, D. M., Bandli, D., & Johnson, S. (2002). Results of salmonella isolation from poultry products, poultry, poultry environment, and other characteristics. *Avian Diseases*, 46(1), 17–24. [https://doi.org/10.1637/0005-2086\(2002\)046\[0017:rosifp\]2.0.co;2](https://doi.org/10.1637/0005-2086(2002)046[0017:rosifp]2.0.co;2)

Sady, S., Błaszczyk, A., Kozak, W., Boryło, P., & Szindler, M. (2021). Quality Assessment of innovative chitosan-based biopolymers for edible food packaging applications. *Food Packaging and Shelf Life*, 30, 100756. <https://doi.org/10.1016/j.fpsl.2021.100756>

Santosh Ku, H.T. et al. (2011) ‘Effects of processing practices on the physico-chemical, microbiological and sensory quality of fresh chicken meat’, *International Journal of Meat Science*, 2(1), pp. 1–6. doi:10.3923/ijmeat.2012.1.6.

Sathishkumar, & Kulkarni, V. V. (2020). Physicochemical and sensory quality of meat in commercial native chicken, backyard native chicken, commercial broiler and spent layer chicken. *Journal of Meat Science*, 15(2), 6–11. <https://doi.org/10.5958/2581-6616.2020.00013.4>

Shen, C., & Zhang, Y. (2023). Evaluate antimicrobial activity of chlorine water on apples and measurement of free chlorine concentrations. *Food Microbiology Laboratory for the Food Science Student*, 81–88. [https://doi.org/10.1007/978-3-031-26197-8\\_11](https://doi.org/10.1007/978-3-031-26197-8_11)

Silva, A. S., Sampaio, A. P., Santos, M. S., Souza, B. W., & Evangelista-Barreto, N. S. (2019). Effect of chitosan coating on contamination of fresh bovine meat sold in The open&nbs; &nbs; &nbs; &nbs; market. *REVISTA CIÊNCIA AGRONÔMICA*, 50(1). <https://doi.org/10.5935/1806-6690.20190005>

Sirri, F. et al. (2017) ‘Effect of EU electrical stunning conditions on breast meat quality of broiler chickens,’ Poultry Science, 96(8), pp. 3000–3004. Available at: <https://doi.org/10.3382/ps/pex048>.

Siriken, B. et al. (2015) ‘Prevalence and characterization of salmonella isolated from chicken meat in Turkey’, Journal of Food Science, 80(5). doi:10.1111/1750-3841.12829.

Škorpilová, T., Šístková, I., Adamcová, M., Pohůnek, V., Kružík, V., & Ševčík, R. (2019). Measuring citrate synthase activity as an enzymatic approach to the differentiation of chilled and frozen/thawed meat. Meat Science, 158, 107856. <https://doi.org/10.1016/j.meatsci.2019.05.027>

Songjiang, Z. and Lixiang, W. (2009) ‘Amyloid-beta associated with chitosan nano-carrier has favorable immunogenicity and permeates the BBB’, AAPS PharmSciTech, 10(3). doi:10.1208/s12249-009-9279-1.

Soultos, N., Tzikas, Z., Abraham, A., Georgantelis, D., & Ambrosiadis, I. (2008). Chitosan effects on quality properties of greek style fresh pork sausages. Meat Science, 80(4), 1150–1156. <https://doi.org/10.1016/j.meatsci.2008.05.008>

Song, D.-H., Ham, Y.-K., Noh, S.-W., Chin, K. B., & Kim, H.-W. (2020). Evaluation of nacl and kcl salting effects on technological properties of pre- and post-rigor chicken breasts at various Ionic strengths. Foods, 9(6), 721. <https://doi.org/10.3390/foods9060721>

Tierrez, A. and García-del Portillo, F. (2005) ‘New concepts in salmonella virulence: The importance of reducing the intracellular growth rate in the host’, Cellular Microbiology, 7(7), pp. 901–909. doi:10.1111/j.1462-5822.2005.00540.x.

TSAI, L.-S., SCHADE, J. E., & MOLYNEUX, B. T. (1992). Chlorination of poultry chiller water: Chlorine demand and disinfection efficiency. Poultry Science, 71(1), 188–196. <https://doi.org/10.3382/ps.0710188>

Van Toan, N., Thi Hanh, T., & Vo Minh Thien, P. (2013). Antibacterial activity of chitosan on some common food contaminating microbes. The Open Biomaterials Journal, 4, 1–5. <https://doi.org/10.2174/1876502501304010001>

Vijayababu, P. et al. (2019) ‘Patulin interference with ATP binding cassette transferring auto inducer –2 in salmonella typhi and biofilm inhibition via quorum sensing’, Informatics in Medicine Unlocked, 16, p. 100230. doi:10.1016/j.imu.2019.100230.

Wabeck, C. J. (2002). Quality assurance and food safety—chicken meat. Commercial Chicken Meat and Egg Production, 871–887. [https://doi.org/10.1007/978-1-4615-0811-3\\_44](https://doi.org/10.1007/978-1-4615-0811-3_44)

Wang, D. et al. (2014) ‘Enzyme-functionalized electrochemical immunosensor based on electrochemically reduced graphene oxide and polyvinyl alcohol-polydimethylsiloxane for the detection of *Salmonella Pullorum* & *Salmonella gallinarum*’, RSC Adv., 4(101), pp. 57733–57742. doi:10.1039/c4ra09901j.

Wessels, K., Rip, D., & Gouws, P. (2021). *Salmonella* in chicken meat: Consumption, outbreaks, characteristics, current control methods and the potential of bacteriophage use. Foods, 10(8), 1742. <https://doi.org/10.3390/foods10081742>

ZHAO, Z.-K. and WANG, Y.-F. (2010) ‘The function of histone chaperones during development’, Hereditas (Beijing), 32(1), pp. 41–48. doi:10.3724/sp.j.1005.2010.00041.

Zheng, L.-Y., & Zhu, J.-F. (2003). Study on antimicrobial activity of chitosan with different molecular weights. Carbohydrate Polymers, 54(4), 527–530. <https://doi.org/10.1016/j.carbpol.2003.07.009>