

## REFERENCES

- Ahmed, J., Shrivhare, U. S., & Raghavan, G. S. V. (2004). Thermal degradation kinetics of anthocyanin and visual colour of plum puree. *European Food Research and Technology*, 218(6), 525-528. doi: 10.1007/s00217-004-0906-5
- Ainsworth, E. A., & Gillespie, K. M. (2007). Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin-Ciocalteu reagent. *Nature Protocols*, 2(4), 875-877. doi: 10.1038/nprot.2007.102
- Amaglianì, L., O'Regan, J., Kelly, A. L., & O'Mahony, J. A. (2017). The composition, extraction, functionality and applications of rice proteins: A review. *Trends in Food Science & Technology*, 64, 1-12. doi: 10.1016/j.tifs.2017.01.008
- Andersen, Ø. M., & Jordheim, M. (2010). Anthocyanins. *Encyclopedia of Life Sciences (eLS)*. doi: 10.1002/9780470015902.a0001909.pub2
- AOAC. (1990). Official methods of analysis of the AOAC, 15th ed. Arlington, VA.
- Benzie, I. F. F., & Strain, J. J. (1996). The Ferric Reducing Ability of Plasma (FRAP) as a Measure of "Antioxidant Power": The FRAP Assay. *Analytical Biochemistry*, 239(1), 70-76. doi: 10.1006/abio.1996.0292
- Boonla, O., Kukongviriyapan, U., Pakdeechote, P., Kukongviriyapan, V., Pannangpatch, P., & Thawornchinsombut, S. (2015). Peptides-Derived from Thai Rice Bran Improves Endothelial Function in 2K-1C Renovascular Hypertensive Rats. *Nutrients*, 7(7), 5783-5799. doi: 10.3390/nu7075252
- Boonloh, K., Kukongviriyapan, V., Kongyinyo, B., Kukongviriyapan, U., Thawornchinsombut, S., & Pannangpatch, P. (2015). Rice Bran Protein Hydrolysates Improve Insulin Resistance and Decrease Pro-inflammatory Cytokine Gene Expression in Rats Fed a High Carbohydrate-High Fat Diet. *Nutrients*, 7(8), 6313-6329. doi: 10.3390/nu7085292
- Boonloh, K., Kukongviriyapan, U., Pannangpatch, P., Kongyinyo, B., Senggunpree, L., Prawan, A., ... & Kukongviriyapan, V. (2015). Rice bran protein hydrolysates prevented interleukin-6- and high glucose-induced insulin resistance in HepG2 cells. *Food & Function*, 6(2), 566-573. doi: 10.1039/C4FO00872C
- Boonloh, K., Lee, E. S., Kim, H. M., Kwon, M. H., Kim, Y. M., Pannangpatch, P., ... & Chung, C. H. (2018). Rice bran protein hydrolysates attenuate diabetic nephropathy in diabetic animal model. *European Journal of Nutrition*, 57(2), 761-772. doi: 10.1007/s00394-016-1366-y
- Branen, A. L., Davidson, P. M., Salminen, S., & Thorngate, J. H. (2001). *Food Additives, Second Edition, Revised and Expanded*. New York, NY: Marcel Dekker.
- Brewer, M. S. (2011). Natural Antioxidants: Sources, Compounds, Mechanisms of Action, and Potential Applications. *Comprehensive Reviews in Food Science and Food Safety*, 10(4), 221-247. doi: 10.1111/j.1541-4337.2011.00156.x
- Brownmiller, C., Howard, L. R., & Prior, R. L. (2008). Processing and Storage Effects on Monomeric Anthocyanins, Percent Polymeric Color, and Antioxidant Capacity of Processed Blueberry Products. *Journal of Food Science*, 73(5), H72-H79. doi: 10.1111/j.1750-3841.2008.00761.x
- Bueno, J. M., Sáez-Plaza, P., Ramos-Escudero, F., Jiménez, A. M., Fett, R., & Asuero, A. G. (2012). Analysis and Antioxidant Capacity of Anthocyanin Pigments. Part II: Chemical Structure, Color, and Intake of Anthocyanins. *Critical Reviews in Analytical Chemistry*, 42(2), 126-151. doi: 10.1080/10408347.2011.632314

- Buvé, C., Kebede, B. T., De Batselier, C., Carrillo, C., Pham, H. T. T., Hendrickx, M., ... & Van Loey, A. (2018). Kinetics of colour changes in pasteurised strawberry juice during storage. *Journal of Food Engineering*, 216, 42-51. doi: 10.1016/j.jfoodeng.2017.08.002
- Carocho, M., & Ferreira, I. C. (2013). A review on antioxidants, prooxidants and related controversy: Natural and synthetic compounds, screening and analysis methodologies and future perspectives. *Food and Chemical Toxicology*, 51, 15-25. doi: 10.1016/j.fct.2012.09.021
- Castañeda-Ovando, A., de Lourdes Pacheco-Hernández, M., Páez-Hernández, M. E., Rodríguez, J. A., & Galán-Vidal, C. A. (2009). Chemical studies of anthocyanins: A review. *Food Chemistry*, 113(4), 859-871. doi: 10.1016/j.foodchem.2008.09.001
- Cheetangdee, N. (2014). Effects of Rice Bran Protein Hydrolysates on the Physicochemical Stability of Oil-in-Water Emulsions. *Journal of Oleo Science*, 63(12), 1231-1241. doi: 10.5650/jos.ess14030
- Chung, C., Rojanasasithara, T., Mutilangi, W., & McClements, D. J. (2015). Enhanced stability of anthocyanin-based color in model beverage systems through whey protein isolate complexation. *Food Research International*, 76, 761-768. doi: 10.1016/j.foodres.2015.07.003
- Cisse, M., Vaillant, F., Kane, A., Ndiaye, O., & Dornier, M. (2012). Impact of the extraction procedure on the kinetics of anthocyanin and colour degradation of roselle extracts during storage. *Journal of the Science of Food and Agriculture*, 92(6), 1214-1221. doi: 10.1002/jsfa.4685
- Cao, S., Liu, L., Lu, Q., Xu, Y., Pan, S., & Wang, K. (2009). Integrated effects of ascorbic acid, flavonoids and sugars on thermal degradation of anthocyanins in blood orange juice. *European Food Research and Technology*, 228(6), 975-983. doi: 10.1007/s00217-009-1015-2
- Danışman, G., Arslan, E., & Toklucu, A. K. (2015). Kinetic Analysis of Anthocyanin Degradation and Polymeric Colour Formation in Grape Juice during Heating. *Czech Journal of Food Sciences*, 33(2), 103-108. doi: 10.17221/446/2014-CJFS
- Fabian, C., & Ju, Y. H. (2011). A Review on Rice Bran Protein: Its Properties and Extraction Methods. *Critical Reviews in Food Science and Nutrition*, 51(9), 816-827. doi: 10.1080/10408398.2010.482678
- Fernandes, I., Marques, C., Évora, A., Faria, A., Calhau, C., Mateus, N., & de Freitas, V. (2018). Anthocyanins: Nutrition and Health. *Bioactive Molecules in Food*, 1-37. doi: 10.1007/978-3-319-54528-8\_79-1
- Food and Agriculture Organization of the United Nations. (2018). Rice Market Monitor. Retrieved from <http://www.fao.org/3/i9243EN/i9243en.pdf>
- Garzón, G. A., & Wrolstad, R. E. (2002). Comparison of the Stability of Pelargonidin-based Anthocyanins in Strawberry Juice and Concentrate. *Journal of Food Science*, 67(4), 1288-1299. doi: 10.1111/j.1365-2621.2002.tb10277.x
- Guan, Y., & Zhong, Q. (2015). The improved thermal stability of anthocyanins at pH 5.0 by gum arabic. *LWT-Food Science and Technology*, 64(2), 706-712. doi: 10.1016/j.lwt.2015.06.018
- Gul, K., Yousuf, B., Singh, A. K., Singh, P., & Wani, A. A. (2015). Rice bran: Nutritional values and its emerging potential for development of functional food—A review. *Bioactive Carbohydrates and Dietary Fibre*, 6(1), 24-30. doi: 10.1016/j.bcdf.2015.06.002
- Hager, A., Howard, L. R., Prior, R. L., & Brownmiller, C. (2008). Processing and Storage Effects on Monomeric Anthocyanins, Percent Polymeric Color, and Antioxidant Capacity of Processed Black Raspberry Products. *Journal of Food Science*, 73(6), H134-H140. doi: 10.1111/j.1750-3841.2008.00855.x

- Hossain, M. A., & Shah, M. D. (2015). A study on the total phenols content and antioxidant activity of essential oil and different solvent extracts of endemic plant Merremia borneensis. *Arabian Journal of Chemistry*, 8(1), 66-71. doi: 10.1016/j.arabjc.2011.01.007
- Hou, Z., Qin, P., Zhang, Y., Cui, S., & Ren, G. (2013). Identification of anthocyanins isolated from black rice (*Oryza sativa L.*) and their degradation kinetics. *Food Research International*, 50(2), 691-697. doi: 10.1016/j.foodres.2011.07.037
- Hu, C., Zawistowski, J., Ling, W., & Kitts, D. D. (2003). Black rice (*Oryza sativa L. indica*) Pigmented Fraction Suppresses both Reactive Oxygen Species and Nitric Oxide in Chemical and Biological Model Systems. *Journal of Agricultural and Food Chemistry*, 51(18), 5271-5277. doi: 10.1021/jf034466n
- Huang, H. T. (1955). Fruit Color Destruction, Decolorization of Anthocyanins by Fungal Enzymes. *Journal of Agricultural and Food Chemistry*, 3(2), 141-146. doi: 10.1021/jf60048a006
- Hubermann, E. M., Heins, A., Stöckmann, H., & Schwarz, K. (2006). Influence of acids, salt, sugars and hydrocolloids on the colour stability of anthocyanin rich black currant and elderberry concentrates. *European Food Research and Technology*, 223(1), 83-90. doi: 10.1007/s00217-005-0139-2
- Jackman, R. L., Yada, R. Y., Tung, M. A., & Speers, R. A. (1987). Anthocyanins as Food Colorants—A Review. *Journal of Food Biochemistry*, 11(3), 201-247. doi: 10.1111/j.1745-4514.1987.tb00123.x
- Jiang, T., Mao, Y., Sui, L., Yang, N., Li, S., Zhu, Z., ... & He, Y. (2019). Degradation of anthocyanins and polymeric color formation during heat treatment of purple sweet potato extract at different pH. *Food Chemistry*, 274, 460-470. doi: 10.1016/j.foodchem.2018.07.141
- Juliano, B. O. (1985). Rice properties and processing. *Food Reviews International*, 1(3), 423-445. doi: 10.1080/87559128509540778
- Kaewjumpol, G., Oruna-Concha, M. J., Niranjan, K., & Thawornchinsombut, S. (2018). The production of hydrolysates from industrially defatted rice bran and its surface image changes during extraction. *Journal of the Science of Food and Agriculture*, 98(9), 3290-3298. doi: 10.1002/jsfa.8832
- Kapcum, N., Uriyapongson, J., Alli, I., & Phimphilai, S. (2016). Anthocyanins, phenolic compounds and antioxidant activities in colored corn cob and colored rice bran. *International Food Research Journal*, 23(6), 2347-2356.
- Kaewjumpol, G., & Thawornchinsombut, S. (2018). Enhanced storage stability of freeze dried rice bran hydrolysates by maltodextrin. *Asia-Pacific Journal of Science and Technology*, 23(2). doi: 10.14456/kkurj.2018.5
- Kim, M. K., Kim, H. A., Koh, K., Kim, H. S., Lee, Y. S., & Kim, Y. H. (2008). Identification and quantification of anthocyanin pigments in colored rice. *Nutrition Research and Practice*, 2(1), 46-49. doi: 10.4162/nrp.2008.2.1.46
- Kırca, A., Özkan, M., & Cemeroğlu, B. (2007). Effects of temperature, solid content and pH on the stability of black carrot anthocyanins. *Food Chemistry*, 101(1), 212-218. doi: 10.1016/j.foodchem.2006.01.019
- Lamuela-Raventós, R. M. (2018). Folin–Ciocalteu method for the measurement of total phenolic content and antioxidant capacity. *Measurement of Antioxidant Activity and Capacity: Recent Trends and Applications*, 107-115. doi: 10.1002/9781119135388.ch6

- Laokuldilok, T., & Rattanathanan, Y. (2014). Protease Treatment for the Stabilization of Rice Bran: Effects on Lipase Activity, Antioxidants, and Lipid Stability. *Cereal Chemistry*, 91(6), 560-565. doi: 10.1094/CCHEM-02-14-0022-R
- Loypimai, P., Moongngarm, A., & Chottanom, P. (2016). Thermal and pH degradation kinetics of anthocyanins in natural food colorant prepared from black rice bran. *Journal of Food Science and Technology*, 53(1), 461-470. doi: 10.1007/s13197-015-2002-1
- Luh, B. S., Barber, S., & de Barber, C. B. (1991). Rice bran: chemistry and technology. In *Rice* (pp. 732-781). Boston, MA: Springer.
- Markakis, P., & Jurd, L. (1974). Anthocyanins and their stability in foods. *Critical Reviews in Food Science & Nutrition*, 4(4), 437-456. doi: 10.1080/10408397409527165
- Martín, J., Navas, M. J., Jiménez-Moreno, A. M., & Asuero, A. G. (2017). Anthocyanin pigments: Importance, sample preparation and extraction. In *Phenolic Compounds-Natural Sources, Importance and Applications*. IntechOpen. doi: 10.5772/66892
- Martynenko, A., & Chen, Y. (2016). Degradation kinetics of total anthocyanins and formation of polymeric color in blueberry hydrothermodynamic (HTD) processing. *Journal of Food Engineering*, 171, 44-51. doi: 10.1016/j.jfoodeng.2015.10.008
- Mateus, N., & de Freitas, V. (2008). Anthocyanins as food colorants. In *Anthocyanins* (pp. 284-304). New York, NY: Springer.
- Miller, N. J., & Rice-Evans, C. A. (1997). Factors Influencing the Antioxidant Activity Determined by the ABTS•+ Radical Cation Assay. *Free Radical Research*, 26(3), 195-199. doi: 10.3109/10715769709097799
- Nayak, B., Berrios, J. D. J., Powers, J. R., & Tang, J. (2011). Thermal Degradation of Anthocyanins from Purple Potato (Cv. Purple Majesty) and Impact on Antioxidant Capacity. *Journal of Agricultural and Food Chemistry*, 59(20), 11040-11049. doi: 10.1021/jf201923a
- Niu, L., Wu, L., & Xiao, J. (2017). Inhibition of gelatinized rice starch retrogradation by rice bran protein hydrolysates. *Carbohydrate Polymers*, 175, 311-319. doi: 10.1016/j.carbpol.2017.07.070
- Park, H. Y., Lee, K. W., & Choi, H. D. (2017). Rice bran constituents: immunomodulatory and therapeutic activities. *Food & Function*, 8(3), 935-943. doi: 10.1039/C6FO01763K
- Pengkumsri, N., Chaiyasut, C., Saenjum, C., Sirilun, S., Peerajan, S., Suwannalert, P., ... & Sivamaruthi, B. S. (2015). Physicochemical and antioxidative properties of black, brown and red rice varieties of northern Thailand. *Food Science and Technology*, 35(2), 331-338. doi: 10.1590/1678-457X.6573
- Phongthai, S., Homthawornchoo, W., & Rawdkuen, S. (2017). Preparation, properties and application of rice bran protein: A review. *International Food Research Journal*, 24(1), 25-34.
- Pisoschi, A. M. & Negulescu, G. P. (2011). Methods for Total Antioxidant Activity Determination: A Review. *Biochemistry & Analytical Biochemistry*, 1(1), 106. doi: 10.4172/2161-1009.1000106
- Poiana, M. A., Alexa, E., & Mateescu, C. (2012). Tracking antioxidant properties and color changes in low-sugar bilberry jam as effect of processing, storage and pectin concentration. *Chemistry Central Journal*, 6(1), 4-14. doi: 10.1186/1752-153X-6-4
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., & Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*, 26(9-10), 1231-1237. doi: 10.1016/S0891-5849(98)00315-3

- Sánchez-Rangel, J. C., Benavides, J., Heredia, J. B., Cisneros-Zevallos, L., & Jacobo-Velázquez, D. A. (2013). The Folin–Ciocalteu assay revisited: improvement of its specificity for total phenolic content determination. *Analytical Methods*, 5(21), 5990-5999. doi: 10.1039/C3AY41125G
- Senaphan, K., Sangartit, W., Pakdeechote, P., Kukongviriyapan, V., Pannangpetch, P., Thawornchinsombut, S., ... & Kukongviriyapan, U. (2018). Rice bran protein hydrolysates reduce arterial stiffening, vascular remodeling and oxidative stress in rats fed a high-carbohydrate and high-fat diet. *European Journal of Nutrition*, 57(1), 219-230. doi: 10.1007/s00394-016-1311-0
- Sharif, M. K., Butt, M. S., Anjum, F. M., & Khan, S. H. (2014). Rice Bran: A Novel Functional Ingredient. *Critical Reviews in Food Science and Nutrition*, 54(6), 807-816. doi: 10.1080/10408398.2011.608586
- Shih, F. F. (2003). An update on the processing of high-protein rice products. *Food/Nahrung*, 47(6), 420-424. doi: 10.1002/food.200390093
- Sivamaruthi, B. S., Kesika, P., & Chaiyasut, C. (2018). Anthocyanins in Thai rice varieties: distribution and pharmacological significance. *International Food Research Journal*, 25(5), 2024-2032.
- Spellman, D., McEvoy, E., O'Cuinn, G., & FitzGerald, R. J. (2003). Proteinase and exopeptidase hydrolysis of whey protein: Comparison of the TNBS, OPA and pH stat methods for quantification of degree of hydrolysis. *International Dairy Journal*, 13(6), 447-453. doi: 10.1016/S0958-6946(03)00053-0
- Supawong, S., Thawornchinsombut, S., & Park, J. W. (2018). Controlling Lipid Oxidation and Volatile Compounds in Frozen Fried Fish Cake Prepared with Rice Bran Hydrolysate. *Journal of Aquatic Food Product Technology*, 27(8), 885-899. doi: 10.1080/10498850.2018.1508103
- Sutharut, J., & Sudarat, J. (2012). Total anthocyanin content and antioxidant activity of germinated colored rice. *International Food Research Journal*, 19(1), 215-221.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L., & Byrne, D. H. (2006). Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis*, 19(6-7), 669-675. doi: 10.1016/j.jfca.2006.01.003
- Tsai, P. J., & Huang, H. P. (2004). Effect of polymerization on the antioxidant capacity of anthocyanins in Roselle. *Food Research International*, 37(4), 313-318. doi: 10.1016/j.foodres.2003.12.007
- Turfan, Ö., Türkyılmaz, M., Yemiş, O., & Özkan, M. (2011). Anthocyanin and colour changes during processing of pomegranate (*Punica granatum* L. cv. Hicaznar) juice from sacs and whole fruit. *Food Chemistry*, 129(4), 1644-1651. doi: 10.1016/j.foodchem.2011.06.024
- Türkyılmaz, M., & Özkan, M. (2012). Kinetics of anthocyanin degradation and polymeric colour formation in black carrot juice concentrates during storage. *International Journal of Food Science & Technology*, 47(11), 2273-2281. doi: 10.1111/j.1365-2621.2012.03098.x
- Wang, W. D., & Xu, S. Y. (2007). Degradation kinetics of anthocyanins in blackberry juice and concentrate. *Journal of Food Engineering*, 82(3), 271-275. doi: 10.1016/j.jfoodeng.2007.01.018
- Wrolstad, R. E. (2004). Anthocyanin Pigments—Bioactivity and Coloring Properties. *Journal of Food Science*, 69(5), C419-C425. doi: 10.1111/j.1365-2621.2004.tb10709.x
- Wrolstad, R. E., Durst, R. W., & Lee, J. (2005). Tracking color and pigment changes in anthocyanin products. *Trends in Food Science & Technology*, 16(9), 423-428. doi: 10.1016/j.tifs.2005.03.019

Wu, J., Guan, Y., & Zhong, Q. (2015). Yeast mannoproteins improve thermal stability of anthocyanins at pH 7.0. *Food Chemistry*, 172, 121-128. doi: 10.1016/j.foodchem.2014.09.059

Yang, Z., Han, Y., Gu, Z., Fan, G., & Chen, Z. (2008). Thermal degradation kinetics of aqueous anthocyanins and visual color of purple corn (*Zea mays L.*) cob. *Innovative Food Science & Emerging Technologies*, 9(3), 341-347. doi: 10.1016/j.ifset.2007.09.001