

References:

- Abraham, E. P., & Chain, E. (1940). An enzyme from bacteria able to destroy penicillin. *Nature*, 146(3713), 837.
- Agaoglu, S., Dostbil, N. & Alemdar, S. (2005). Antimicrobial effect of seed extract of cardamom (ElettariacardamomumMaton). *Yyu Vet FakDerg*. 16(2): 99-101.
- Ahn, B.-Z., Yoon, Y.-D., Lee, Y.H., Kim, B.-H., Sok, D.-E., 1998. In-hibitory effect of bupleuri radix saponins on adhesion of some solid tumor cells and relation to hemolytic action: Screening of 232 herbal drugs for anti-cell adhesion. *Planta Medica* 64, 220–224
- Akinyemi, K. O., Oluwa, O. K., & Omomigbehin, E. O. (2006). Antimicrobial activity of crude extracts of three medicinal plants used in south-west Nigerian folk medicine on some food borne bacterial pathogens. *African Journal of Traditional, Complementary and Alternative Medicines*, 3(4), 13-22.
- Aladesanmi, A. J., Oriola, A. O., Oguntimehin, S. A., Akinkunmi, E. O., Igbeneghu, O. A., & Obuotor, E. M. (2019). Comparative antimicrobial and antioxidant activities of four medicinal plants. *Ife Journal of Science*, 21(2), 59-66.
- Al Laham, S. A. & MOHAMMAD, F. (2014). Screening the Antimicrobial Activity of some Medicinal Plants against Multidrug Resistance Escherichia Coli Type (1). *Asian Journal of Pharmaceutical Research and Health Care*, 6(1).
- Arabski, M., Węgierek-Ciuk, A., Czerwonka, G., Lankoff, A. & Kaca, W. (2012). Effects of saponins against clinical Escherichia coli strains and eukaryotic cell line. *BioMed Research International*, 2012.
- Azwanida, N. N. (2015). A review on the extraction methods use in medicinal plants, principle, strength and limitation. *Med Aromat Plants*, 4(196), 2167-0412.
- Balouiri, M., Sadiki, M., & Ibnsouda, S. K. (2016). Methods for in vitro evaluating antimicrobial activity: A review. *Journal of pharmaceutical analysis*, 6(2), 71-79.
- Bankole, O. M. (2018). Phytochemical and antimicrobial studies of colloidal silver nanoparticles mediated by Laporteaestuans extract. *International Journal of Biomedical Engineering and Clinical Science*, 4(2), 58.
- Baumann, E., Stoya, G., Völkner, A., Richter, W., Lemke, C., Linss, W., 2000. Hemolysis of human erythrocytes with saponin affects the membrane structure. *Acta Histochemica* 102, 21–35
- Berg, H. C. (2008). *Escherichia coli in Motion*. Springer Science & Business Media.
- Beuria, T. K., Santra, M. K., & Panda, D. (2005). Sanguinarine blocks cytokinesis in bacteria by inhibiting FtsZ assembly and bundling. *Biochemistry*, 44(50), 16584-16593.
- Bradford, P. A. (2001). Extended-spectrum β -lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clinical microbiology reviews*, 14(4), 933-951.

Brink, M. (2019). Laportea aestuans (PROTA) - PlantUse English. Retrieved from [https://uses.plantnet-project.org/en/Laportea_aestuans_\(PROTA\)](https://uses.plantnet-project.org/en/Laportea_aestuans_(PROTA))

Bruneton, J. 1995. Pharmacognosy, Phytochemistry, Medicinal Plants. Lavoisier Publishing, Paris, pp. 538–544 (ISBN 2-4730-0028-7).

Capasso, L. (1998). 5300 years ago, the Ice Man used natural laxatives and antibiotics. *The Lancet*, 352(9143), 1864.

Cheesman, M. J., Ilanko, A., Blonk, B., & Cock, I. E. (2017). Developing new antimicrobial therapies: are synergistic combinations of plant extracts/compounds with conventional antibiotics the solution?. *Pharmacognosy reviews*, 11(22), 57.

Chin, F. S., Chong, K. P., Markus, A., & Wong, N. K. (2013). Tea polyphenols and alkaloids content using soxhlet and direct extraction methods. *World J Agric Sci*, 9(3), 266-270.

Cohen, M. B. & Giannella, R. A. (1992). Hemorrhagic colitis associated with Escherichia coli O157:H7. *Advances in internal medicine*, 37, 173–195.

Ćujić, N., Šavikin, K., Janković, T., Pljevljakušić, D., Zdunić, G., & Ibrić, S. (2016). Optimization of polyphenols extraction from dried chokeberry using maceration as traditional technique. *Food chemistry*, 194, 135–142. <https://doi.org/10.1016/j.foodchem.2015.08.008>

Davis, W. W., & Stout, T. R. (1971). Disc Plate Method of Microbiological Antibiotic Assay: I. Factors Influencing Variability and Error. *Appl. Environ. Microbiol.*, 22(4), 659-665.

Demarque, D.P., Callejon, D.R., deOliveira, G.G., Silva, D.B, Carollo, C.A., & Lopes, N.P. (2018) The role of tannins as antiulcer agents: a fluorescence-imaging based study. *Revista Brasileira de Farmacognosia*, 28 (4), 425-432

European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID). (2003). Determination of minimum inhibitory concentrations (MICs) of antibacterial agents by broth dilution. *Clinical Microbiology and Infection*, 9(8), ix-xv.

Farkas, A., Maróti, G., Kereszt, A. & Kondorosi, É. (2017). Comparative analysis of the bacterial membrane disruption effect of two natural plant antimicrobial peptides. *Frontiers in microbiology*, 8, 51.
Field. J. A., Kortekaas, S. and Lettinga. G. (1989) *Biol. Wastes* 29. 241.

Ginovyan, M., Petrosyan, M. & Trchounian, A. (2017). Antimicrobial activity of some plant materials used in Armenian traditional medicine. *BMC complementary and alternative medicine*, 17(1), 50.

Godowski, K. C. (1989). Antimicrobial action of sanguinarine. *The Journal of clinical dentistry*, 1(4), 96-101.

Goh, Z-H. (2016, May 22) What's the reaction between phenol and ferric chloride? Retrieved from <https://www.quora.com/Whats-the-reaction-between-phenol-and-ferric-chloride>

Hamoud, R., Reichling, J. & Wink, M. (2014). Synergistic antimicrobial activity of combinations of sanguinarine and EDTA with vancomycin against multidrug-resistant bacteria. *Drug metabolism letters*, 8(2), 119-128

Harborne, J. B., & Williams, C. A. (2000). Advances in flavonoid research since 1992. *Phytochemistry*, 55(6), 481-504.

Haslam, E. (1989) *Plant Polyphenols*. Cambridge University Press, Cambridge.

Hertsenberg, A. & Noori, M. (2010, February). Plate Streaking & Growth.

Holla, G., Yeluri, R., & Munshi, A. K. (2012). Evaluation of minimum inhibitory and minimum bactericidal concentration of nano-silver base inorganic anti-microbial agent (Novaron®) against streptococcus mutans. *Contemporary clinical dentistry*, 3(3), 288.

Hooper, D. C. (2001). Emerging mechanisms of fluoroquinolone resistance. *Emerging infectious diseases*, 7(2), 337.

Hosking, G. A. (2001). *Russia and the Russians: A history*. Harvard University Press.

Hertsenberg, A., & Noori, M. (2010, February 12). *protocol:plate streaking and growth*. Retrieved from http://fg.cns.utexas.edu/fg/protocol%3A_plate_streaking_and_growth.html

Hudzicki, J. (2009). Kirby-Bauer disk diffusion susceptibility test protocol.

Iorizzi, M., Lanzotti, V., Ranalli, G., De Marino, S., Zollo, F., 2002. Antimicrobial furostanol saponins from the seeds of Capsicum annuum L. var.acuminatum. *Journal of Agricultural and Food Chemistry* 50,4310–4316

Itabashi, M., Segawa, K., Ikeda, Y., Kondo, S., Naganawa, H., Koyano, T., Umezawa, K., 1999. A new bioactive steroidal saponin, furcrestatin,from the plant Furcraea foetida. *Carbohydrate Research* 323, 57–62.

Jorgensen, J. H., & Turnidge, J. D. (2007). Susceptibility test methods: dilution and disk diffusion methods, p. 1152–1172. In P. R. Murray, E. J. Baron, J. H. Jorgensen, M. L. Landry, and M. A. Pfaller (ed.), *Manual of clinical microbiology*, 9th ed.

Jovanović, A. A., Đorđević, V. B., Zdunić, G. M., Pljevljakušić, D. S., Šavikin, K. P., Gođevac, D. M., & Bugarski, B. M. (2017). Optimization of the extraction process of polyphenols from Thymus serpyllum L. herb using maceration, heat-and ultrasound-assisted techniques. *Separation and Purification Technology*, 179, 369-380.

Just, M.J., Recio, M.C., Giner, R.M., Cuéllar, M.J., Mañez, S., Bilia,A.R., R'ios, J.-L., 1998. Anti-inflammatory activity of unusual lupane saponins from Bupleurum fruticosescens. *Planta Medica* 64, 404–407.

Karth, E. S., Ranjitha, P. & Mohankumar, A. (2009). Antimicrobial Potential of Plant Seed Extracts against Multidrug-resistant Methicillin Resistant *Staphylococcus aureus* (MDR-MRSA). International Journal of Biology.

Khan, A., Haque, E., Rahman, M. & Nessa, F. (2008). Bioactivity of roots of *Laporteacrenulata*. Pharmaceutical Biology, 46(10-11), 695-699.

Lambert, R. J. W., Skandamis, P. N., Coote, P. J., & Nychas, G. J. (2001). A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. Journal of applied microbiology, 91(3), 453-462.

Li, P., Xu, G., Li, S. P., Wang, Y. T., Fan, T. P., Zhao, Q. S. & Zhang, Q. W. (2008). Optimizing ultraperformance liquid chromatographic analysis of 10 diterpenoid compounds in *Salvia miltiorrhiza* using central composite design. Journal of agricultural and food chemistry, 56(4), 1164–1171.
<https://doi.org/10.1021/jf073020u>

Mambe, F. T., Voukeng, I. K., Beng, V. P. & Kuete, V. (2016). Antibacterial activities of methanol extracts from *Alchorneacordifolia* and four other Cameroonian plants against MDR phenotypes. Journal of Taibah University Medical Sciences, 11(2), 121-127.

Mergaert, P., Nikovics, K., Kelemen, Z., Maunoury, N., Vaubert, D., Kondorosi, A. & Kondorosi, E. (2003). A novel family in *Medicago truncatula* consisting of more than 300 nodule-specific genes coding for small, secreted polypeptides with conserved cysteine motifs. Plant physiology, 132(1), 161–173.
<https://doi.org/10.1104/pp.102.018192>

Moher, D., Liberati, A., Tetzlaff, J. & Altman, D. G. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097.
<https://doi.org/10.1371/journal.pmed.1000097>

Morrison, J. F. (2012). Comparative studies on the in vitro antioxidant and antibacterial activity of methanolic and hydro-ethanolic extracts from eight edible leafy vegetables of Ghana (Doctoral dissertation).

Murray, P. R., Baron, E. J., Jorgensen, J. H., Landry, M. L. & Pfaller, M. A. (2007). Manual of Clinical Microbiology. Ed. 9th.

Nascimento, G. G., Locatelli, J., Freitas, P. C. & Silva, G. L. (2000). Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. Brazilian journal of microbiology, 31(4), 247-256.

Nath, M. C., Chakravorty, M. K. & Chowdhury, S. R. (1946). Liebermann-Burchard reaction for steroids. Nature, 157(3978), 103.

Newman, D. J., Cragg, G. M. & Snader, K. M. (2000). The influence of natural products upon drug discovery. Natural product reports, 17(3), 215-234.

Nishizawa, K., Nakata, I., Kishida, A., Ayer, W. A. and Browne, L. M. (1990) Phytochemistry 29, 2491. Oeding P. (1992). Koleraen i Bergen 1848-49--hvordan oppstod epidemien? [Cholera in Bergen in 1848-1849--what caused the epidemic?]. Tidsskrift for den Norske laegeforening :tidsskrift for praktiskmedisin, nyraekke, 112(17), 2214–2217.

Oloyede, G. K., & Ayanbadejo, O. E. (2014). Phytochemical, toxicity, antimicrobial and antioxidant screening of extracts obtained from Laporteaestuans (Gaud). *J Med Sci*, 14(51), e59.

Oloyede, G. K., & Oyelola, M. S. (2013). Chrysen-2-ol derivative from West Indian Wood Nettle Laporteaestuans (L.) Chew inhibits oxidation and microbial growth in vitro. *EXCLI journal*, 12, 894–906.

Oloyede, G. K. (2016). Toxicity, antimicrobial and antioxidant activities of methyl salicylate dominated essential oils of Laporteaestuans (Gaud). *Arabian Journal of Chemistry*, 9, S840-S845.

Oloyede, G. K. & Ayanbadejo, O. E. (2014). Phytochemical, toxicity, antimicrobial and antioxidant screening of extracts obtained from Laporteaestuans (Gaud). *J Med Sci*, 14(51), e59.

Oyekunle, M. A., Aiyelaagbe, O. O., & Fafunso, M. A. (2006). Evaluation of the antimicrobial activity of saponins extract of Sorghum bicolor L. Moench. *African journal of Biotechnology*, 5(23).

Pretorius, C. J. & Watt, E. (2001). Purification and identification of active components of Carpobrotus edulis L. *J. Ethnopharmacol.* 76: 87-91.

Rahal, J. J., & Simberkoff, M. S. (1979). Bactericidal and bacteriostatic action of chloramphenicol against meningeal pathogens. *Antimicrobial agents and chemotherapy*, 16(1), 13-18.

Rahman, M. A., Rana, M. S., Zaman, M. M., Uddin, S. A. & Akter, R. (2010). Antioxidant, antibacterial and cytotoxic activity of the methanol extract of UrticaCrenulata. *Journal of Scientific Research*, 2(1), 169-177.

Rahman, M. M., Khan, A., Haque, M. E., & Rahman, M. M. (2008). Antimicrobial and cytotoxic activities of Laporteacrenulata. *Fitoterapia*, 79(7-8), 584-586.

Rasigade, J. P. & Vandenesch, F. (2014). *Staphylococcus aureus*: a pathogen with still unresolved issues. *Infection, Genetics and Evolution*, 21, 510-514.

Rates, S. M. K. (2001). Plants as source of drugs. *Toxicon*, 39(5), 603-613.

Rehse, K., and Kawerau, G. (1974). Mechanism of the reaction of aromatic compounds with formaldehyde in concentrated sulfuric acid (Marquis reagent). *Archiv der Pharmazie (Weinheim)*, 307, 934-942.

Roussel, C., Cordonnier, C., Livrelli, V., Van de Wiele, T. & Blanquet-Diot, S. (2017). Enterotoxigenic and enterohemorrhagic Escherichia coli: Survival and modulation of virulence in the human gastrointestinal tract.

Samie, A. (Ed.). (2017). *Escherichia coli: Recent Advances on Physiology, Pathogenesis and Biotechnological Applications*. BoD–Books on Demand.

Samuel, Y. (2019, Feb 2). *How does Salkowski's Test for Steroid content work?* Retrieved from <https://www.researchgate.net/post/how does Salkowskis Test for Steroid content work>

Sangi, M.S., Momuat, L.I. dan Kumaunang, M., 2013. Uji toksitas dan skriningfitokimiatepunggabahpelelaharen (Arangepinnata). Manado: Universitas Sam Ratulangi.

- Scalbert, A. (1991). Antimicrobial properties of tannins. *Phytochemistry*, 30(12), 3875–3883.
- Schouten, M. A., Willems, R. J., Kraak, W. A., Top, J., Hoogkamp-Korstanje, J. A. & Voss, A. (2001). Molecular analysis of Tn1546-like elements in vancomycin-resistant enterococci isolated from patients in Europe shows geographic transposon type clustering. *Antimicrobial agents and chemotherapy*, 45(3), 986-989.
- Schroeder, C. M., Meng, J., Zhao, S., DebRoy, C., Torcolini, J., Zhao, C.,...& White, D. G. (2002). Antimicrobial resistance of *Escherichia coli* O26, O103, O111, O128, and O145 from animals and humans. *Emerging infectious diseases*, 8(12), 1409.
- Schroeder, C. M., Zhao, C., DebRoy, C., Torcolini, J., Zhao, S., White, D. G.,...& Meng, J. (2002). Antimicrobial resistance of *Escherichia coli* O157 isolated from humans, cattle, swine, and food. *Applied and Environmental Microbiology*, 68(2), 576-581.
- Silva, N. C. C. & Fernandes Júnior, A. (2010). Biological properties of medicinal plants: a review of their antimicrobial activity. *Journal of venomous Animals and Toxins including tropical diseases*, 16(3), 402-413.
- Simaremare, E. (2014). SKRINING FITOKIMIA EKSTRAK ETANOL DAUN GATAL (Laporteadeicumana (Roxb.) Wedd
- Simaremare, E. S., Holle, E., & Budi, M. (2015). Perbandingan Efektifitas Antinyeri Salep Daun Gatal Dari Simplicia Laporteadeicumana dan Laportea sp. *PHARMACY: Jurnal Farmasi Indonesia (Pharmaceutical Journal of Indonesia)*, 12(1), 1-10.
- Simaremare, E. S., Ruban, A. & Runtuboi, D. Y. (2017). Aktivitas antibakteri ekstrak ketanoldaungatal (Laportea aestivalis (L.) Chew). *Jurnal Biologi Papua*, 9(1), 1-7.
- Simaremare, E. S., Gunawan, E., Yarangga, I., Satya, M. D., & Yabansabra, Y. R. (2020, July). Antibacterial and Toxicity Activities Itchy Leaves (Laporteadeicumana, Roxb. Wedd) Extract. In *Journal of Physics: Conference Series* (Vol. 1503, No. 1, p. 012041). IOP Publishing.
- Simões, C.M.O., Amoros, M., Girre, L., 1999. Mechanism of antiviral activity of triterpenoid saponins. *Phytotherapy Research* 13, 323–328.
- Sonfack, G., FossiTchinda, C., Simo, I. K., Bitchagno, G. T. M., Nganou, B. K., Çelik, İ.,...& Kuete, V. (2019). Saponin with antibacterial activity from the roots of *Albizia adianthifolia*. *Natural product research*, 1-9.
- Sparg, S., Light, M. E. & Van Staden, J. (2004). Biological activities and distribution of plant saponins. *Journal of ethnopharmacology*, 94(2-3), 219-243.
- Stojanović, G., Radulović, N., Hashimoto, T. & Palić, R. (2005). In vitro antimicrobial activity of extracts of four *Achillea* species: The composition of *Achillea clavennae* L. (Asteraceae) extract. *Journal of ethnopharmacology*, 101(1-3), 185-190.

Tagousop, C. N., Kengne, I. C., Ngnokam, D. & Voutquenne-Nazabadioko, L. (2018). Antimicrobial activities of saponins from *Melanthera elliptica* and their synergistic effects with antibiotics against pathogenic phenotypes. *Chemistry Central Journal*, 12(1), 97.

Taylor, T. A. & Unakal, C. G. (2017). *Staphylococcus aureus*.

Tepal, P. (2016). Phytochemical screening, total flavonoid and phenolic content assays of various solvent extracts of tepal of *Musa paradisiaca*. *Malaysian Journal of Analytical Sciences*, 20(5), 1181-1190.

Traore, F., Faure, R., Olivier, E., Gasquet, M., Azas, N., Debrauwer, L., Keita, A., Timon-David, P., Balansard, G., 2000. Structure and antiprotozoal activity of triterpenoid saponins from *Glinus oppositifolius*. *Planta Medica* 66, 368–371.

Trieu-Cuot, P., G. De Cespedes, F. Bentorcha, F. Delbos, E. Gaspar, and T. Horaud. 1993. Study of heterogeneity of chloramphenicol acetyltransferase (CAT) genes in streptococci and enterococci by polymerase chain reaction: Characterization of a new CAT determinant. *Antimicrob. Agents Chemother.* 37:2593–2598.

Trusheva B, Trunkova D and Bankova V. (2008) Different extraction methods of biologically active components from propolis: a preliminary study. *Chem Cen J*, 2007; 1: 13.

Tyler, V.E., Brady, L.R., Robbers, J.E., 1981. *Pharmacognosy*, eighth ed. Lea&Ferbiger, Philadelphia, p. 67 (ISBN 0-8121-0793-4).

Ultee, A., Bennik, M. H. J. & Moezelaar, R. J. A. E. M. (2002). The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Applied and environmental microbiology*, 68(4), 1561-1568.

Urs, N. V. R. R. and Dunleavy. J. M. (1975) *Phytopathology* 65, 686.

Vineetha, N., Vignesh, R. A. & Sridhar, D. (2015). Preparation, Standardization of Antibiotic Discs and Study of Resistance Pattern for First-Line Antibiotics in Isolates from Clinical Samples. *International Journal of Applied Research*, 1(11), 624-631.

Wintola, O. A. & Afolayan, A. J. (2015). The antibacterial, phytochemicals and antioxidants evaluation of the root extracts of *Hydnora africana* Thunb. used as antidysenteric in Eastern Cape Province, South Africa. *BMC complementary and alternative medicine*, 15(1), 307.

Wright, G. D. 1999. Aminoglycoside-modifying enzymes. *Curr. Opin. Microbiol.* 2:499–503.

Xu, F. X., Yuan, C., Wan, J. B., Yan, R., Hu, H., Li, S. P., & Zhang, Q. W. (2015). A novel strategy for rapid quantification of 20(S)-protopanaxatriol and 20(S)-protopanaxadiol saponins in *Panax notoginseng* P. ginseng and *P. quinquefolium*. *Natural product research*, 29(1), 46–52.

<https://doi.org/10.1080/14786419.2014.957698>

Yeo, Y. L., Chia, Y. Y., Lee, C. H., Sow, H. S., & Yap, W. S. (2014). Effectiveness of maceration periods with different extraction solvents on in-vitro antimicrobial activity from fruit of *Momordica charantia* L. *Journal of Applied Pharmaceutical Science*, 4(10), 016-020.

- Zaffiri, L., Gardner, J., & Toledo-Pereyra, L. H. (2012). History of antibiotics. From salvarsan to cephalosporins. *Journal of Investigative Surgery*, 25(2), 67-77.
- Zhang, Q. W., Lin, L. G., & Ye, W. C. (2018). Techniques for extraction and isolation of natural products: a comprehensive review. *Chinese medicine*, 13(1), 20.
- Zhang, Y., Kensler, T. W., Cho, C. G., Posner, G. H. & Talalay, P. (1994). Anticarcinogenic activities of sulforaphane and structurally related synthetic norbornyl isothiocyanates. *Proceedings of the National Academy of Sciences*, 91(8), 3147-3150.
- Zoecklein B (2006) Tannins, color and the 2006 season, continued: phenolic compounds and red wine processing, *Enology Notes* #117. Retrieved from
<http://www.apps.fst.vt.edu/extension/enology/EN/117.html>
- Zypler, A., Słomińska, M., & Namieśnik, J. (2012). 2.04-Soxhlet Extraction and New Developments Such as Soxtec. *Comprehensive Sampling and Sample Preparation*, 65-82.