

## CHAPTER VII – REFERENCES

- Aerts, O., & Goossens, A. (2021). Contact Allergy to Preservatives. In J. D. Johansen, V. Mahler, J.-P. Lepoittevin, & P. J. Frosch (Eds.), *Contact Dermatitis* (pp. 835–876). Springer International Publishing. [https://doi.org/10.1007/978-3-030-36335-2\\_85](https://doi.org/10.1007/978-3-030-36335-2_85)
- Almeida, A., Sarmento, B., & Rodrigues, F. (2017). Insights on in vitro models for safety and toxicity assessment of cosmetic ingredients. *International Journal of Pharmaceutics*, 519(1–2), 178–185. <https://doi.org/10.1016/j.ijpharm.2017.01.024>
- Amorim, S., Reis, C. A., Reis, R. L., & Pires, R. A. (2021). Extracellular matrix mimics using hyaluronan-based biomaterials. *Trends in Biotechnology*, 39(1), 90–104. <https://doi.org/10.1016/j.tibtech.2020.06.003>
- Arda, O., Göksügür, N., & Tüzün, Y. (2014). Basic histological structure and functions of facial skin. *Clinics in Dermatology*, 32(1), 3–13. <https://doi.org/10.1016/j.cldermatol.2013.05.021>
- Armengot-Carbo, M., Hernández-Martín, Á., & Torrelo, A. (2015). The role of filaggrin in the skin barrier and disease development. *Actas Dermo-Sifiliográficas (English Edition)*, 106(2), 86–95. <https://doi.org/10.1016/j.adengl.2014.12.007>
- Atolani, O., Adamu, N., Oguntoye, O. S., Zubair, M. F., Fabiyi, O. A., Oyegoke, R. A., Adeyemi, O. S., Areh, E. T., Tarigha, D. E., Kambizi, L., & Olantuji, G. A. (2020). Chemical characterization, antioxidant, cytotoxicity, Anti-Toxoplasma gondii and antimicrobial potentials of the Citrus sinensis seed oil for sustainable cosmeceutical production. *Helix*. [https://www.cell.com/helix/pdf/S2405-8440\(20\)30244-9.pdf](https://www.cell.com/helix/pdf/S2405-8440(20)30244-9.pdf)
- Avantaggiato, A., Bertuzzi, G., Vitiello, U., Iannucci, G., Pasin, M., Pascali, M., Cervelli, V., & Carinci, F. (2014). Role of antioxidants in dermal aging: An in vitro study by Q-RT-PCR. *Aesthetic Plastic Surgery*, 38(5), 1011–1016. <https://doi.org/10.1007/s00266-014-0380-9>
- Barco, D., & Giménez-Arnau, A. (2008). Xerosis: A dysfunction of the epidermal barrier. *Actas Dermo-Sifiliográficas (English Edition)*, 99(9), 671–682. [https://doi.org/10.1016/s1578-2190\(08\)70343-3](https://doi.org/10.1016/s1578-2190(08)70343-3)
- Barnes, T. M., Mijaljica, D., Townley, J. P., Spada, F., & Harrison, I. P. (2021). Vehicles for drug delivery and cosmetic moisturizers: Review and comparison. *Pharmaceutics*, 13(12), 2012. <https://doi.org/10.3390/pharmaceutics13122012>
- Baroni, A., Buommino, E., De Gregorio, V., Ruocco, E., Ruocco, V., & Wolf, R. (2012). Structure and function of the epidermis related to barrier properties. *Clinics in Dermatology*, 30(3), 257–262. <https://doi.org/10.1016/j.cldermatol.2011.08.007>
- Berardesca, E., Farage, M., & Maibach, H. (2012). Sensitive skin: An overview. *International Journal of Cosmetic Science*, 35(1), 2–8. <https://doi.org/10.1111/j.1468-2494.2012.00754.x>
- Bhardwaj, M., Masand, N., Sahoo, J., & Patil, V. M. (2020). Risk Assessment of Cosmetic Preservatives Using QSAR: QSAR of Cosmetic Preservatives. *International Journal of Quantitative Structure-Property Relationships*, 5(1), 44–62. <https://doi.org/10.4018/IJQSPR.2020010103>
- Bitschar, K., Wolz, C., Krismer, B., Peschel, A., & Schittek, B. (2017). Keratinocytes as sensors and central players in the immune defense against *Staphylococcus aureus* in the skin. *Journal of Dermatological Science*, 87(3), 215–220. <https://doi.org/10.1016/j.jdermsci.2017.06.003>
- Boer, D. E. C., van Smeden, J., Al-Khakany, H., Melnik, E., van Dijk, R., Absalah, S., Vreeken, R. J., Haenen, C. C. P., Lavrijsen, A. P. M., Overkleef, H. S., Aerts, J. M. F. G., & Bouwstra, J. A. (2020). Skin of atopic dermatitis patients shows disturbed  $\beta$ -glucocerebrosidase and acid

sphingomyelinase activity that relates to changes in stratum corneum lipid composition. *Biochimica Et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, 1865(6), 158673. <https://doi.org/10.1016/j.bbalip.2020.158673>

Boer, M., Duchnik, E., Maleszka, R., & Marchlewicz, M. (2016). Structural and biophysical characteristics of human skin in maintaining proper epidermal barrier function. *Advances in Dermatology and Allergology*, 1, 1–5. <https://doi.org/10.5114/pdia.2015.48037>

Cha, H. J., He, C., Zhao, H., Dong, Y., An, I.-S., & An, S. (2016). Intercellular and intracellular functions of ceramides and their metabolites in skin (review). *International Journal of Molecular Medicine*, 38(1), 16–22. <https://doi.org/10.3892/ijmm.2016.2600>

Choi, H. K., Cho, Y. H., Lee, E. O., Kim, J. W., & Park, C. S. (2017). Phytosphingosine enhances moisture level in human skin barrier through stimulation of the filaggrin biosynthesis and degradation leading to NMF formation. *Archives of Dermatological Research*, 309(10), 795–803. <https://doi.org/10.1007/s00403-017-1782-8>

Colella, G., Vicedomini, A., Soro, V., Lanza, A., & Cirillo, N. (2012). Molecular insights into the effects of sodium hyaluronate preparations in keratinocytes. *Clinical and Experimental Dermatology*, 37(5), 516–520. <https://doi.org/10.1111/j.1365-2230.2011.04279.x>

de Koning, H. D., van den Bogaard, E. H., Bergboer, J. G. M., Kamsteeg, M., van Vlijmen-Willems, I. M. J. J., Hitomi, K., Henry, J., Simon, M., Takashita, N., Ishida-Yamamoto, A., Schalkwijk, J., & Zeeuwen, P. L. J. M. (2012). Expression profile of cornified envelope structural proteins and keratinocyte differentiation-regulating proteins during skin barrier repair. *British Journal of Dermatology*, 166(6), 1245–1254. <https://doi.org/10.1111/j.1365-2133.2012.10885.x>

Draelos, Z. D. (2012). New treatments for restoring impaired epidermal barrier permeability: Skin barrier repair creams. *Clinics in Dermatology*, 30(3), 345–348. <https://doi.org/10.1016/j.clindermatol.2011.08.018>

Duarte, I., Silveira, J. E., Hafner, M. de, Toyota, R., & Pedroso, D. M. (2017). Sensitive skin: Review of an ascending concept. *Anais Brasileiros De Dermatologia*, 92(4), 521–525. <https://doi.org/10.1590/abd1806-4841.201756111>

Eckl, K.-M., Tidhar, R., Thiele, H., Oji, V., Hausser, I., Brodesser, S., Preil, M.-L., Önal-Akan, A., Stock, F., Müller, D., Becker, K., Casper, R., Nürnberg, G., Altmüller, J., Nürnberg, P., Traupe, H., Futerman, A. H., & Hennies, H. C. (2013). Impaired epidermal ceramide synthesis causes autosomal recessive congenital ichthyosis and reveals the importance of ceramide acyl chain length. *Journal of Investigative Dermatology*, 133(9), 2202–2211. <https://doi.org/10.1038/jid.2013.153>

Evrard, C., Lambert de Rouvroit, C., & Poumay, Y. (2021). Epidermal hyaluronan in barrier alteration-related disease. *Cells*, 10(11), 3096. <https://doi.org/10.3390/cells10113096>

Fallacara, A., Baldini, E., Manfredini, S., & Vertuani, S. (2018). Hyaluronic acid in the Third Millennium. *Polymers*, 10(7), 701. <https://doi.org/10.3390/polym10070701>

Gendrisch, F., Nováčková, A., Sochorová, M., Haarhaus, B., Vávrová, K., Schempp, C. M., & Wölflle, U. (2020). Gentiana lutea Extract Modulates Ceramide Synthesis in Primary and Psoriasis-Like Keratinocytes. *Molecules*, 25(8), 1832. <https://doi.org/10.3390/molecules25081832>

Grether-Beck, S., Felsner, I., Brenden, H., Kohne, Z., Majora, M., & Marini, A. et al. (2012). Urea Uptake Enhances Barrier Function and Antimicrobial Defense in Humans by Regulating Epidermal Gene Expression. *Journal Of Investigative Dermatology*, 132(6), 1561-1572. doi: 10.1038/jid.2012.42

- Gruber, J. V., Holtz, R., & Riemer, J. (2021). Hyaluronic acid (HA) stimulates the in vitro expression of CD44 proteins but not HAS1 proteins in normal human epidermal keratinocytes (nheks) and is HA molecular weight dependent. *Journal of Cosmetic Dermatology*. <https://doi.org/10.1111/jocd.14188>
- Guéniche, A., Bastien, P., Ovigne, J. M., Kermici, M., Courchay, G., Chevalier, V., Breton, L., & Castiel-Higounenc, I. (2009). Bifidobacterium Longum Lysate, a new ingredient for reactive skin. *Experimental Dermatology*, 19(8). <https://doi.org/10.1111/j.1600-0625.2009.00932.x>
- Habeebuddin, M., Karnati, R. K., Shiroorkar, P. N., Nagaraja, S., Asdaq, S. M. B., Khalid Anwer, Md., & Fattepur, S. (2022). Topical Probiotics: More Than a Skin Deep. *Pharmaceutics*, 14(3), 557. <https://doi.org/10.3390/pharmaceutics14030557>
- Hamilton, D. F., Ghert, M., & Simpson, A. H. R. W. (2015). Interpreting regression models in clinical outcome studies. *Bone & Joint Research*, 4(9), 152–153. <https://doi.org/10.1302/2046-3758.49.2000571>
- Hänel, K., Cornelissen, C., Lüscher, B., & Baron, J. (2013). Cytokines and the skin barrier. *International Journal of Molecular Sciences*, 14(4), 6720–6745. <https://doi.org/10.3390/ijms14046720>
- Hänel, K. H., Pfaff, C. M., Cornelissen, C., Amann, P. M., Marquardt, Y., Czaja, K., Kim, A., Lüscher, B., & Baron, J. M. (2016). Control of the physical and antimicrobial skin barrier by an IL-31–IL-1 signaling network. *The Journal of Immunology*, 196(8), 3233–3244. <https://doi.org/10.4049/jimmunol.1402943>
- Hanyu, O., Nakae, H., Miida, T., Higashi, Y., Fuda, H., & Endo, M. et al. (2012). Cholesterol sulfate induces expression of the skin barrier protein filaggrin in normal human epidermal keratinocytes through induction of ROR $\alpha$ . *Biochemical And Biophysical Research Communications*, 428(1), 99–104. doi: 10.1016/j.bbrc.2012.10.013
- Hartini, H., Vlorensia, Abdullah, H., Martinus, A. R., & Ikhtiar, R. (2019). The effect of a moisturizing cream containing saccharide isomerase and ceramide on reducing transepidermal water loss in eczema. *Proceedings of the International Conference on Health Informatics and Medical Application Technology*. <https://doi.org/10.5220/0009516204110417>
- Hirabayashi, T., Anjo, T., Kaneko, A., Senoo, Y., Shibata, A., Takama, H., Yokoyama, K., Nishito, Y., Ono, T., Taya, C., Muramatsu, K., Fukami, K., Muñoz-Garcia, A., Brash, A. R., Ikeda, K., Arita, M., Akiyama, M., & Murakami, M. (2017). PNPLA1 has a crucial role in skin barrier function by directing acylceramide biosynthesis. *Nature Communications*, 8(1). <https://doi.org/10.1038/ncomms14609>
- Hon, K. L., Leung, A. K., & Barankin, B. (2013). Barrier repair therapy in atopic dermatitis: An overview. *American Journal of Clinical Dermatology*, 14(5), 389–399. <https://doi.org/10.1007/s40257-013-0033-9>
- How, K. N., Yap, W. H., Lim, C. L., Goh, B. H., & Lai, Z. W. (2020). Hyaluronic acid-mediated drug delivery system targeting for inflammatory skin diseases: A Mini Review. *Frontiers in Pharmacology*, 11. <https://doi.org/10.3389/fphar.2020.01105>
- Hsu, M.-F., & Chiang, B.-H. (2009). Stimulating effects of bacillus subtilis natto-fermented radix astragali on hyaluronic acid production in human skin cells. *Journal of Ethnopharmacology*, 125(3), 474–481. <https://doi.org/10.1016/j.jep.2009.07.011>
- Ito, S., Ishikawa, J., Naoe, A., Yoshida, H., Hachiya, A., Fujimura, T., Kitahara, T., & Takema, Y. (2017). Ceramide synthase 4 is highly expressed in involved skin of patients with atopic dermatitis. *Journal*

- of the European Academy of Dermatology and Venereology, 31(1), 135–141. <https://doi.org/10.1111/jdv.13777>
- Jain, A. K., Singh, D., Dubey, K., Maurya, R., Mittal, S., & Pandey, A. K. (2018). Models and Methods for In Vitro Toxicity. In *In Vitro Toxicology* (pp. 45–65). Elsevier. <https://doi.org/10.1016/B978-0-12-804667-8.00003-1>
- Jakasa, I., Thyssen, J. P., & Kezic, S. (2018). The role of skin barrier in occupational contact dermatitis. *Experimental Dermatology*, 27(8), 909–914. <https://doi.org/10.1111/exd.13704>
- Jang, M., Baek, S., Kang, G., Yang, H., Kim, S., & Jung, H. (2020). Dissolving Microneedle with high molecular weight hyaluronic acid to improve skin wrinkles, dermal density and elasticity. *International Journal of Cosmetic Science*, 42(3), 302–309. doi:10.1111/ics.12617
- Jegasothy, S. M., Zabolotniaia, V., & Bielfeldt, S. (2014). Efficacy of a New Topical Nano-hyaluronic Acid in Humans. *The Journal of Clinical and Aesthetic Dermatology*, 7(3), 27–29.
- Jennemann, R., Rabionet, M., Gorgas, K., Epstein, S., Dalpke, A., Rothermel, U., Bayerle, A., van der Hoeven, F., Imgrund, S., Kirsch, J., Nickel, W., Willecke, K., Riezman, H., Gröne, H.-J., & Sandhoff, R. (2011). Loss of ceramide synthase 3 causes lethal skin barrier disruption. *Human Molecular Genetics*, 21(3), 586–608. <https://doi.org/10.1093/hmg/ddr494>
- Jepps, O. G., Dancik, Y., Anissimov, Y. G., & Roberts, M. S. (2013). Modeling the human skin barrier — towards a better understanding of dermal absorption. *Advanced Drug Delivery Reviews*, 65(2), 152–168. <https://doi.org/10.1016/j.addr.2012.04.003>
- Kaewsanit, T., Chakkavittumrong, P., & Waranuch, N. (2021). Clinical Comparison of Topical 2.5% Benzoyl Peroxide plus 5% Niacinamide to 2.5% Benzoyl Peroxide Alone in the Treatment of Mild to Moderate Facial Acne Vulgaris. *Journal of Clinical and Aesthetic Dermatology*, 14(6), 35–41.
- Kahraman, E., Kaykin, M., Şahin Bektay, H., & Güngör, S. (2019). Recent advances on topical application of ceramides to restore barrier function of skin. *Cosmetics*, 6(3), 52. <https://doi.org/10.3390/cosmetics6030052>
- Kezic, S., & Jakasa, I. (2016). Filaggrin and skin barrier function. *Current Problems in Dermatology*, 49, 1–7. <https://doi.org/10.1159/000441539>
- Kim, B. E., & Leung, D. Y. M. (2018). Significance of skin barrier dysfunction in atopic dermatitis. *Allergy, Asthma & Immunology Research*, 10(3), 207. <https://doi.org/10.4168/aair.2018.10.3.207>
- Kim, E., Hwang, K., Lee, J., Han, S., Kim, E., Park, J., & Cho, J. (2018). Skin Protective Effect of Epigallocatechin Gallate. *International Journal Of Molecular Sciences*, 19(1), 173. doi:10.3390/ijms19010173
- Kim, T. H., Kim, W. J., Park, S. Y., Kim, H., & Chung, D. K. (2021). In vitro anti-wrinkle and skin-moisturizing effects of evening primrose (*Oenothera biennis*) sprout and identification of its active components. *Processes*, 9(1), 145. <https://doi.org/10.3390/pr9010145>
- Kim, Y., & Lim, K.-M. (2021). Skin barrier dysfunction and filaggrin. *Archives of Pharmacal Research*, 44(1), 36–48. <https://doi.org/10.1007/s12272-021-01305-x>
- Laviad, E. L., Albee, L., Pankova-Kholmyansky, I., Epstein, S., Park, H., Merrill, A. H., & Futerman, A. H. (2008). Characterization of ceramide synthase 2. *Journal of Biological Chemistry*, 283(9), 5677–5684. <https://doi.org/10.1074/jbc.m707386200>

- Lee, J.-O., Hwang, S.-H., Shen, T., Kim, J. H., You, L., Hu, W., & Cho, J. Y. (2021). Enhancement of skin barrier and hydration-related molecules by protopanaxatriol in human keratinocytes. *Journal of Ginseng Research*, 45(2), 354–360. <https://doi.org/10.1016/j.jgr.2020.12.003>
- Lee, S. E., & Lee, S. H. (2018). Skin barrier and calcium. *Annals of Dermatology*, 30(3), 265. <https://doi.org/10.5021/ad.2018.30.3.265>
- Lodén, M. (2012). Effect of moisturizers on epidermal barrier function. *Clinics in Dermatology*, 30(3), 286–296. <https://doi.org/10.1016/j.cldermatol.2011.08.015>
- Loden, M. (2016). Treatments improving skin barrier function. *Current Problems in Dermatology*, 49, 112–122. <https://doi.org/10.1159/000441586>
- LONG, C. C., & FINLAY, A. Y. (1991). The finger-tip unit—a new practical measure. *Clinical and Experimental Dermatology*, 16(6), 444–447. <https://doi.org/10.1111/j.1365-2230.1991.tb01232.x>
- Love, J., Selker, R., Marsman, M., Jamil, T., Dropmann, D., Verhagen, J., Ly, A., Gronau, Q. F., Smíra, M., Epskamp, S., Matzke, D., Wild, A., Knight, P., Rouder, J. N., Morey, R. D., & Wagenamakers, E.-J. (2019). JASP: Graphical statistical software for common statistical designs. *Journal of Statistical Software*, 88(2). <https://doi.org/10.18637/jss.v088.i02>
- Malaisse, J., Bourguignon, V., De Vuyst, E., Lambert de Rouvroit, C., Nikkels, A. F., Flamion, B., & Poumay, Y. (2014). Hyaluronan metabolism in human keratinocytes and atopic dermatitis skin is driven by a balance of hyaluronan synthases 1 and 3. *Journal of Investigative Dermatology*, 134(8), 2174–2182. <https://doi.org/10.1038/jid.2014.147>
- Martin, E., Zhang, A., & Campiche, R. (2022). Saccharide isomerate ameliorates cosmetic scalp conditions in a Chinese study population. *Journal of Cosmetic Dermatology*, n/a(n/a). <https://doi.org/10.1111/jocd.14913>
- Meckfessel, M. H., & Brandt, S. (2014). The structure, function, and importance of ceramides in skin and their use as therapeutic agents in skin-care products. *Journal of the American Academy of Dermatology*, 71(1), 177–184. <https://doi.org/10.1016/j.jaad.2014.01.891>
- Milani, M., & Sparavigna, A. (2017). The 24-hour skin hydration and barrier function effects of a hyaluronic 1%, glycerin 5%, and centella asiatica stem cells extract moisturizing fluid: An intra-subject, randomized, assessor-blinded study. *Clinical, Cosmetic and Investigational Dermatology*, Volume 10, 311–315. <https://doi.org/10.2147/ccid.s144180>
- Misery, L., Loser, K., & Ständer, S. (2016). Sensitive skin. *Journal of the European Academy of Dermatology and Venereology*, 30, 2–8. <https://doi.org/10.1111/jdv.13532>
- Mizutani, Y., Kihara, A., & Igarashi, Y. (2006). LASS3 (longevity assurance homologue 3) is a mainly testis-specific (dihydro)ceramide synthase with relatively broad substrate specificity. *Biochemical Journal*, 398(Pt 3), 531–538. <https://doi.org/10.1042/BJ20060379>
- Mori, M., Hamamoto, A., Takahashi, A., Nakano, M., Wakikawa, N., Tachibana, S., Ikebara, T., Nakaya, Y., Akutagawa, M., & Kinouchi, Y. (2007). Development of a new water sterilization device with a 365&nbsp;nm UV-led. *Medical & Biological Engineering & Computing*, 45(12), 1237–1241. <https://doi.org/10.1007/s11517-007-0263-1>
- Moss, G. P., Gullick, D. R., & Wilkinson, S. C. (2015). Skin structure and physiology. *Predictive Methods in Percutaneous Absorption*, 1–24. [https://doi.org/10.1007/978-3-662-47371-9\\_1](https://doi.org/10.1007/978-3-662-47371-9_1)
- Mussi, L., Baby, A. R., Camargo Junior, F. B., Padovani, G., Sufi, B. da S., & Magalhães, W. V. (2021). Propanediol (and) Caprylic Acid (and) Xylitol as a New Single Topical Active Ingredient against

Acne: In Vitro and In Vivo Efficacy Assays. *Molecules*, 26(21), 6704. <https://doi.org/10.3390/molecules26216704>

Nashchekina, Y. A., & Raydan, M. (2017). Noninvasive penetration of 5 nm hyaluronic acid molecules across the epidermal barrier (in vitro) and its interaction with human skin cells. *Skin Research and Technology*, 24(1), 129–134. <https://doi.org/10.1111/srt.12400>

Niehues, H., Bouwstra, J. A., El Ghalbzouri, A., Brandner, J. M., Zeeuwen, P. L., & van den Bogaard, E. H. (2018). 3D skin models for 3R research: The potential of 3D reconstructed skin models to study skin barrier function. *Experimental Dermatology*, 27(5), 501–511. <https://doi.org/10.1111/exd.13531>

Panzuti, P., Vidémont, E., Fantini, O., Fardouet, L., Noël, G., Cappelle, J., & Pin, D. (2020). A moisturizer formulated with glycerol and propylene glycol accelerates the recovery of skin barrier function after experimental disruption in dogs. *Veterinary Dermatology*, 31(5), 344. <https://doi.org/10.1111/vde.12859>

Papakonstantinou, E., Roth, M., & Karakiulakis, G. (2012). Hyaluronic acid: A key molecule in skin aging. *Dermato-Endocrinology*, 4(3), 253–258. <https://doi.org/10.4161/derm.21923>

Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research*, 29(9). <https://doi.org/10.1093/nar/29.9.e45>

Proksch, E. (2018). Ph in nature, humans and skin. *The Journal of Dermatology*, 45(9), 1044–1052. <https://doi.org/10.1111/1346-8138.14489>

Puebla-Barragan, S., & Reid, G. (2021). Probiotics in cosmetic and personal care products: Trends and challenges. *Molecules*, 26(5), 1249. <https://doi.org/10.3390/molecules26051249>

Pullmannová, P., Staňková, K., Pospíšilová, M., Školová, B., Zbytovská, J., & Vávrová, K. (2014). Effects of sphingomyelin/ceramide ratio on the permeability and microstructure of model stratum corneum lipid membranes. *Biochimica Et Biophysica Acta (BBA) - Biomembranes*, 1838(8), 2115–2126. <https://doi.org/10.1016/j.bbamem.2014.05.001>

Rabionet, M., Gorgas, K., & Sandhoff, R. (2014). Ceramide synthesis in the epidermis. *Biochimica Et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, 1841(3), 422–434. <https://doi.org/10.1016/j.bbalip.2013.08.011>

Richters, R., Falcone, D., Uzunbajakava, N., Verkruyse, W., van Erp, P., & van de Kerkhof, P. (2014). What is sensitive skin? A systematic literature review of objective measurements. *Skin Pharmacology and Physiology*, 28(2), 75–83. <https://doi.org/10.1159/000363149>

Sahoo, M. R., Varma, S. R., Patel, N. K., Sivaprakasam, T. O., Mishra, A., Muguli, G., Babu, S. K., Babu, U. V., Paramesh, R., & Jadhav, A. N. (2017). In vitro anti-inflammatory and skin protective properties of extract comprising ceramides from de-oiled rice bran: Sahoo et al. *Journal of Food Biochemistry*, 41(3), e12353. <https://doi.org/10.1111/jfbc.12353>

Shin, K.-O., Kim, S., Park, B. D., Uchida, Y., & Park, K. (2021). N-Palmitoyl Serinol Stimulates Ceramide Production through a CB1-Dependent Mechanism in In Vitro Model of Skin Inflammation. *International Journal of Molecular Sciences*, 22(15), 8302. <https://doi.org/10.3390/ijms22158302>

Song, U., & Kim, J. (2020). Assessment of the potential risk of 1,2-hexanediol using phytotoxicity and cytotoxicity testing. *Ecotoxicology and Environmental Safety*, 201, 110796. <https://doi.org/10.1016/j.ecoenv.2020.110796>

- Svec, D., Tichopad, A., Novosadova, V., Pfaffl, M. W., & Kubista, M. (2015). How good is a PCR efficiency estimate: Recommendations for precise and robust qPCR efficiency assessments. *Biomolecular Detection and Quantification*, 3, 9–16. <https://doi.org/10.1016/j.bdq.2015.01.005>
- Szöllősi, A., Guéniche, A., Jammayrac, O., Szabó-Papp, J., Blanchard, C., Vasas, N., András, M., Juhasz, I., Breton, L., & Bíró, T. (2016). Bifidobacterium longum extract exerts pro-differentiating effects on human epidermal keratinocytes, *in vitro*. *Experimental Dermatology*, 26. <https://doi.org/10.1111/exd.13130>
- Takeda, S., Shimoda, H., Takarada, T., & Imokawa, G. (2018). Strawberry seed extract and its major component, tiliroside, promote ceramide synthesis in the stratum corneum of human epidermal equivalents. *PLOS ONE*, 13(10), e0205061. <https://doi.org/10.1371/journal.pone.0205061>
- Takeda, S., Terazawa, S., Shimoda, H., & Imokawa, G. (2021).  $\beta$ -Sitosterol 3-O-D-glucoside increases ceramide levels in the stratum corneum via the up-regulated expression of ceramide synthase-3 and glucosylceramide synthase in a reconstructed human epidermal keratinization model. *PLOS ONE*, 16(3), e0248150. doi: 10.1371/journal.pone.0248150
- Tasić-Kostov, M., & Savić, S. (2015). New Natural Sugar-Based Surfactants Intended for Stabilization of Cosmetic/Dermopharmaceutical Vehicles – Safety and Efficacy Assessment. *Acta Medica Mediana*. <http://scindeks-clanci.ceon.rs/data/pdf/0365-4478/2015/0365-44781501034T.pdf>
- Tsakok, T., Woolf, R., Smith, C. H., Weidinger, S., & Flohr, C. (2018). Atopic dermatitis: The skin barrier and beyond. *British Journal of Dermatology*, 180(3), 464–474. <https://doi.org/10.1111/bjd.16934>
- Varma, S. R., Sivaprakasam, T. O., Arumugam, I., Dilip, N., Raghuraman, M., Pavan, K. B., Rafiq, M., & Paramesh, R. (2019). In vitro anti-inflammatory and skin protective properties of Virgin coconut oil. *Journal of Traditional and Complementary Medicine*, 9(1), 5–14. <https://doi.org/10.1016/j.jtcme.2017.06.012>
- Vávrová, K., Kováčik, A., & Opálka, L. (2017). Ceramides in the Skin Barrier. *European Pharmaceutical Journal*, 64(2), 28–35. <https://doi.org/10.1515/afpuc-2017-0004>
- Vinardell, M. P., & Mitjans, M. (2017). Alternative Methods to Animal Testing for the Safety Evaluation of Cosmetic Ingredients: An Overview. *Cosmetics*, 4(3), 30. <https://doi.org/10.3390/cosmetics4030030>
- Walters, R. M., Mao, G., Gunn, E. T., & Hornby, S. (2012). Cleansing formulations that respect skin barrier integrity. *Dermatology Research and Practice*, 2012, 1–9. <https://doi.org/10.1155/2012/495917>
- Wang, G., Wang, G., Zhang, X., Wang, F., & Song, R. (2011). Isolation of high quality RNA from cereal seeds containing high levels of starch. *Phytochemical Analysis*, 23(2), 159–163. <https://doi.org/10.1002/pca.1337>
- Yamamoto, M., Sassa, T., Kyono, Y., Uemura, H., Kugo, M., Hayashi, H., Imai, Y., Yamanishi, K., & Kihara, A. (2021). Comprehensive stratum corneum ceramide profiling reveals reduced acylceramides in ichthyosis patient with cers3 mutations. *The Journal of Dermatology*, 48(4), 447–456. <https://doi.org/10.1111/1346-8138.15725>
- Yatagai, T., Shimauchi, T., Yamaguchi, H., Sakabe, J.-ichi, Aoshima, M., Ikeya, S., Tatsuno, K., Fujiyama, T., Ito, T., Ojima, T., & Tokura, Y. (2018). Sensitive skin is highly frequent in extrinsic atopic dermatitis and correlates with disease severity markers but not necessarily with skin barrier impairment. *Journal of Dermatological Science*, 89(1), 33–39. <https://doi.org/10.1016/j.jdermsci.2017.10.011>

- Yatsuhashi, H., Furuyashiki, T., Vo, P. H., Kamasaka, H., & Kuriki, T. (2021). Effects of glycogen on ceramide production in cultured human keratinocytes via acid sphingomyelinase activation. *Journal of Applied Glycoscience*, 68(2), 41–46. [https://doi.org/10.5458/jag.jag-2020\\_0012](https://doi.org/10.5458/jag.jag-2020_0012)
- Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., & Madden, T. L. (2012). Primer-Blast: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*, 13(1). <https://doi.org/10.1186/1471-2105-13-134>
- Youn, Y. H. (2020). Enhancing the effect of aronia extract on hyaluronic acid synthesis through liposome formation. *Journal of People, Plants, and Environment*, 23(4), 465–473. <https://doi.org/10.11628/ksppe.2020.23.4.465>
- Zaniboni, M. C., Samorano, L. P., Orfali, R. L., & Aoki, V. (2016). Skin barrier in atopic dermatitis: Beyond filaggrin. *Anais Brasileiros De Dermatologia*, 91(4), 472–478. <https://doi.org/10.1590/abd1806-4841.20164412>
- Zhang, S., Ding, S., Yu, J., Chen, X., Lei, Q., & Fang, W. (2015). Antibacterial Activity, in Vitro Cytotoxicity, and Cell Cycle Arrest of Gemini Quaternary Ammonium Surfactants. *Langmuir*, 31(44), 12161–12169. <https://doi.org/10.1021/acs.langmuir.5b01430>