

REFERENCES

- Abedini, Colin, M., Hubert, J., Charpentier, E., Angelis, A., Bounasri, H., Bertaux, B., Kotland, A., Reffuveille, F., Nuzillard, J.-M., Renault, J.-H., & Gangloff, S. C. (2020). Abundant Extractable Metabolites from Temperate Tree Barks: The Specific Antimicrobial Activity of *Prunus Avium* Extracts. *Antibiotics*, *9*(3), 111. <https://doi.org/10.3390/antibiotics9030111>
- Abubakar, A. R., & Haque, M. (2020). Preparation of Medicinal Plants: Basic Extraction and Fractionation Procedures for Experimental Purposes. *Journal of pharmacy & bioallied sciences*, *12*(1), 1–10. https://doi.org/10.4103/jpbs.JPBS_175_19
- Abreu, A., McBain, A., & Simões, M. (2012). Plants as sources of new antimicrobials and resistance-modifying agents. *Natural Product Reports*, *29*(9), 1007. doi: 10.1039/c2np20035j
- Adebayo, E. A., Ishola, O. R., Taiwo, O. S., Majolagbe, O. N., & Adekeye, B. T. (2009). Evaluations of the methanol extract of *Ficus exasperata* stem bark, leaf and root for phytochemical analysis and antimicrobial activities. *African Journal of Plant Science*, *3*(12), 283–287.
- Akhtar, N., Ihsan-ul-Haq, & Mirza, B. (2018). Phytochemical analysis and comprehensive evaluation of antimicrobial and antioxidant properties of 61 medicinal plant species. *Arabian Journal Of Chemistry*, *11*(8), 1223–1235. doi: 10.1016/j.arabjc.2015.01.013
- Allocati, N., Masulli, M., Alexeyev, M. F., & Di Ilio, C. (2013). *Escherichia coli* in Europe: an overview. *International journal of environmental research and public health*, *10*(12), 6235–6254. <https://doi.org/10.3390/ijerph10126235>
- Aniszewski, T. (2015). *Alkaloids: Chemistry, Biology, Ecology, and Applications*. Netherlands: Elsevier Science.
- Ara, I., Shinwari, M., Rashed, S., & Bakir, M. (2013). Evaluation of Antimicrobial Properties of Two Different Extracts of *Juglans regia* Tree Bark and Search for Their Compounds Using Gas Chromatography-Mass Spectrum. *International Journal Of Biology*, *5*(2), 92–102. doi: 10.5539/ijb.v5n2p92
- Arora, D., & Sood, H. (2017). In vitro antimicrobial potential of extracts and phytoconstituents from *Gymnema sylvestre* R.Br. leaves and their biosafety evaluation. *AMB Express*, *7*(1). doi: 10.1186/s13568-017-0416-z
- Aspé, E., & Fernández, K. (2011). The effect of different extraction techniques on extraction yield, total phenolic, and anti-radical capacity of extracts from *Pinus radiata* Bark. *Industrial Crops And Products*, *34*(1), 838–844. doi: 10.1016/j.indcrop.2011.02.002
- Beattie, A. J., Hay, M., Magnusson, B., de Nys, R., Smeathers, J., & Vincent, J. F. (2011). Ecology and bioprospecting. *Austral ecology*, *36*(3), 341–356. <https://doi.org/10.1111/j.1442->

9993.2010.02170.x

- Belay, R., & Makonnen, E. (2020). Anti-inflammatory activities of ethanol leaves extract and solvent fractions of *Zehneria scabra* (Cucurbitaceae) in rodents. *Asian Journal of Natural Product Biochemistry*, 18(1).
- Blondeau, D, St-Pierre, A, Bourdeau, N, Bley, J, Lajeunesse, A, Desgagné-Penix, I. 2020. Antimicrobial activity and chemical composition of white birch (*Betula papyrifera* Marshall) bark extracts. *MicrobiologyOpen*. 9:e944. <https://doi.org/10.1002/mbo3.944>
- Borges, A., José, H., Homem, V., & Simões, M. (2020). Comparison of Techniques and Solvents on the Antimicrobial and Antioxidant Potential of Extracts from *Acacia dealbata* and *Olea europaea*. *Antibiotics (Basel, Switzerland)*, 9(2), 48. <https://doi.org/10.3390/antibiotics9020048>
- Brennan, M., Fritsch, C., Cosgun, S. et al. (2020). Quantitative and qualitative composition of bark polyphenols changes longitudinally with bark maturity in *Abies alba* Mill.. *Annals of Forest Science* 77, 9. <https://doi.org/10.1007/s13595-019-0916-x>
- Breijyeh, Z., Jubeh, B., & Karaman, R. (2020). Resistance of Gram-Negative Bacteria to Current Antibacterial Agents and Approaches to Resolve It. *Molecules*. 16;25(6):1340. doi: 10.3390/molecules25061340. PMID: 32187986; PMCID: PMC7144564.
- [CDC] Centers for Disease Control and Prevention. (2022). *E. coli (Escherichia coli)*. Retrieved 25 May 2022, from <https://www.cdc.gov/ecoli/index.html>
- Chandrasekaran, V. (2007). Typical Rubber Testing Methods. *Essential Rubber Formulary*, 16-22. doi: 10.1016/b978-081551539-5.50008-0
- Choudary, D.K., Kumar, M., Prasad, R., & Kumar, V. (2018). *In Silico Approach for Sustainable Agriculture*. Germany: Springer Singapore.
- Cobellis, G., Trabalza-Marinucci, M., & Yu, Z. (2016). Critical evaluation of essential oils as rumen modifiers in ruminant nutrition: A review. *Science Of The Total Environment*, 545-546, 556-568. doi: 10.1016/j.scitotenv.2015.12.103
- Cullinane, A., Markey, B., Maguire, D., Leonard, F., & Archambault, M. (2013). *Clinical Veterinary Microbiology E-Book*. United Kingdom: Elsevier Health Sciences.
- Cuong, D., Hoan, N., Dong, D., Thuy, L., Thanh, N., & Ha, H. et al. (2020). Tannins: Extraction from Plants. *Intechopen*, (4), 57-76. doi: 10.5772/intechopen.860.40
- Cushnie, T. P., & Lamb, A. J. (2005). Antimicrobial activity of flavonoids. *International journal of antimicrobial agents*, 26(5), 343–356. <https://doi.org/10.1016/j.ijantimicag.2005.09.002>
- Cushnie, T. P., Cushnie, B., & Lamb, A. J. (2014). Alkaloids: an overview of their antibacterial, antibiotic-enhancing and antivirulence activities. *International journal of antimicrobial agents*, 44(5), 377–386. <https://doi.org/10.1016/j.ijantimicag.2014.06.001>

- De, N., & Ifeoma, E. (2002). Antimicrobial effects of components of the bark extract of neem (*Azadirachta indica* A. Juss). *TECHNOLOGY AND DEVELOPMENT*, 8, 23-28.
- de Boer, J. (2005). POLYCHLORINATED BIPHENYLS. *Encyclopedia Of Analytical Science*, 214-225. doi: 10.1016/b0-12-369397-7/00472-6
- Dewick, P. M. (2009). *Medicinal Natural Products* (3rd ed., p. 311). Chichester: John Wiley & Sons.
- Dhanani, T., Shah, S., Gajbhiye, N., & Kumar, S. (2017). Effect of extraction methods on yield, phytochemical constituents and antioxidant activity of *Withania somnifera*. *Arabian Journal Of Chemistry*, 10, S1193-S1199. doi: 10.1016/j.arabjc.2013.02.015
- Dhayalan, A., Gracilla, D. E., Dela Peña, R. A., Jr, Malison, M. T., & Pangilinan, C. R. (2018). Phytochemical Constituents and Antimicrobial Activity of the Ethanol and Chloroform Crude Leaf Extracts of *Spathiphyllum cannifolium* (Dryand. ex Sims) Schott. *Journal of pharmacy & bioallied sciences*, 10(1), 15–20. https://doi.org/10.4103/jpbs.JPBS_95_17
- Ejikeme, C., Ezeonu, C. S., & Eboatu, A. N. (2014). Determination of Physical and Phytochemical Constituents of some Tropical Timbers Indigenous to nigerdelta area of nigeria. *European Scientific Journal*, 10(18), 247-270.
- Ene, A.C., Atawodi, S.E., Ameh, DA, Kwanashie, HO & Agomo, Philip. (2009). In vivo antiplasmodial effect of chloroform extracts of *Artemisia maciverae* Linn and *Artemisia maritima* Linn. *AFRICAN JOURNAL OF BIOTECHNOLOGY*. 8. 10.4314/ajb.v8i23.66362.
- Ezeonu, C. S., & Ejikeme, C. M. (2016). Qualitative and quantitative determination of phytochemical contents of indigenous Nigerian softwoods. *New Journal of Science*, 2016.
- Fu, J., Wu, Z., & Zhang, L. (2019). Clinical applications of the naturally occurring or synthetic glycosylated low molecular weight drugs. *Progress In Molecular Biology And Translational Science*, 487-522. doi: 10.1016/bs.pmbts.2019.03.005
- Giri, Dil & Rajbhandari, Meena. (2018). PHYTOCHEMICAL ANALYSIS AND CONSTITUENTS OF HEXANE EXTRACT OF MELASTOMA MALABATHRICUM L. *Journal of Institute of Science and Technology*. 23. 18-25. 10.3126/jist.v23i1.22150.
- Goodwin, A. C., Steele-Moore, L., & Schwalbe, R. (2007). *Antimicrobial Susceptibility Testing Protocols*. United States: CRC Press.
- Hazimah, H., Azharman, Z., Yuharmen, Y., Rahyuti, V., & Afriliani, A. (2019). Antibacterial Activity of The Hexane Extract from Leaf of Plant *Solanum Ferox* L. *Proceedings Of The Proceedings Of The 5Th Annual International Seminar On Trends In Science And Science Education, AISTSSE 2018, 18-19 October 2018, Medan, Indonesia*. doi: 10.4108/eai.18-10-2018.2287176
- Hudzicki, J. (2009). Kirby-Bauer disk diffusion susceptibility test protocol. *American society for microbiology*, 15, 55-63.

- Hon, D D.-S, & Shiraishi, N. (2001). *Wood and Cellulosic Chemistry, Second Edition, Revised, and Expanded*. Italy: Taylor & Francis.
- Houghton, P., & Raman, A. (2012). *Laboratory handbook for the fractionation of natural extracts*. Springer Science & Business Media.
- Jabran, K. (2019). *Role of Mulching in Pest Management and Agricultural Sustainability*. Germany: Springer International Publishing.
- Julianto, T. S. (2019). *Fitokimia: Tinjauan Metabolit Sekunder dan Skrining Fitokimia*. Yogyakarta: Universitas Islam Indonesia.
- Juwitaningsih, T., Sari, S., Jahro, I., Silaban, S., & Simorangkir, M. (2021). Study of Phytochemical, Antibacterial Activity and Toxicity on Acetone Extract Seed *LeersiaHexandraSw*. *Journal Of Physics: Conference Series*, 1811(1), 012130. doi: 10.1088/1742-6596/1811/1/012130
- Jyske, T., Brännström, H., Halmemies, E., Laakso, T., Kilpeläinen, P., & Hyvönen, J. et al. (2022). Stilbenoids of Norway spruce bark: does the variability caused by raw-material processing offset the biological variability?. *Biomass Conversion And Biorefinery*. doi: 10.1007/s13399-022-02624-9
- Kataoka, H. (2018). Pharmaceutical Analysis | Sample Preparation ☆ *Reference Module In Chemistry, Molecular Sciences And Chemical Engineering*. doi: 10.1016/b978-0-12-409547-2.14358-6
- Karunaratne, D., & Pamunuwa, G. (2017). *Food Additives*. Croatia: IntechOpen.
- Khalifa, M. A., & Al Sayed Omar, O. (2022). *High Performance Liquid Chromatography: Theory, Instrumentation and Application in Drug Quality Control*. Germany: De Gruyter.
- Khameneh, B., Iranshahy, M., Soheili, V., & Fazly Bazzaz, B. (2019). Review on plant antimicrobials: a mechanistic viewpoint. *Antimicrobial Resistance & Infection Control*, 8(1). doi: 10.1186/s13756-019-0559-6
- Khoddami, A., Wilkes, M.A., & Roberts, T.H. (2013). Techniques for Analysis of Plant Phenolic Compounds. *Molecules*, 18(2):2328-2375. <https://doi.org/10.3390/molecules18022328>
- Kuete, V., Fozing, D., Kapche, W., Mbaveng, A., Kuate, J., Ngadjui, B., & Abegaz, B. (2009). Antimicrobial activity of the methanolic extract and compounds from *Morus mesozygia* stem bark. *Journal Of Ethnopharmacology*, 124(3), 551-555. doi: 10.1016/j.jep.2009.05.004
- Kumar, S., Prasad, S. R., Banerjee, R., Agarwal, D. K., Kulkarni, K. S., & Ramesh, K. V. (2017). Green solvents and technologies for oil extraction from oilseeds. *Chemistry Central journal*, 11, 9. <https://doi.org/10.1186/s13065-017-0238-8>
- Kwieceński, J. M., & Horswill, A. R. (2020). Staphylococcus aureus bloodstream infections: pathogenesis and regulatory mechanisms. *Current opinion in microbiology*, 53, 51–60. <https://doi.org/10.1016/j.mib.2020.02.005>

- Lense, O.Ī. (2012). The wild plants used as traditional medicines by indigenous people of Manokwari, West Papua. *Biodiversitas*, 13, 98-106.
- Loree, J., & Lappin, S. (2021). Bacteriostatic Antibiotics. Retrieved 29 May 2022, from <https://www.ncbi.nlm.nih.gov/books/NBK547678/>
- Madivoli, E. S., Maina, E. G., Kairigo, P. K., Murigi, M. K., Ogilo, J. K., Nyangau, J. O., Kimani, P. K., Kipyegon, C. (2018). In vitro antioxidant and antimicrobial activity of *Prunus africana* (Hook. f.) Kalkman (bark extracts) and *Harrisonia abyssinica* Oliv. extracts (bark extracts): A comparative study. *Journal of Medicinal Plants for Economic Development*, 2(1), –. doi:10.4102/jomped.v2i1.39
- Mahizan, N. A., Yang, S. K., Moo, C. L., Song, A. A., Chong, C. M., Chong, C. W., Abushelaibi, A., Lim, S. E., & Lai, K. S. (2019). Terpene Derivatives as a Potential Agent against Antimicrobial Resistance (AMR) Pathogens. *Molecules (Basel, Switzerland)*, 24(14), 2631. <https://doi.org/10.3390/molecules24142631>
- Makut, M., Gyar, S., Pennap, G., & Anthony, P. (2008). Phytochemical screening and antimicrobial activity of the ethanolic and methanolic extracts of the leaf and bark of *Khaya senegalensis*. *African Journal Of Biotechnology*, 7(9), 1216-1219.
- Mallikharjuna, P. B., & Seetharam, Y. N. (2009). *In Vitro* Antimicrobial Screening of Alkaloid Fractions from *Strychnos potatorum*. *E-Journal Of Chemistry*, 6(4), 1200–1204.
- Manunta, P., Ferrandi, M., & Bianchi, G. (2004). Ouabain. *Encyclopedia Of Endocrine Diseases*, 447-450. doi: 10.1016/b0-12-475570-4/00957-4
- Martins, A., Salgueiro, L., Gonçalves, M., Vila, R., Tomi, F., & Adzet, T. et al. (2000). Antimicrobial Activity and Chemical Composition of the Bark Oil of *Croton stellulifer*, an Endemic Species from S. Tomé e Príncipe. *Planta Medica*, 66(7), 647-650. doi: 10.1055/s-2000-8623
- Masabni, J. (n.d.). Mulching - Should you water before or after mulching?. Retrieved 22 May 2022, from <https://agrillifeextension.tamu.edu/library/gardening/mulching/#:~:text=Mulching%20is%20a%20long%2Destablished,to%20keep%20the%20fruit%20clean.>
- McMaster, H. (2015). What is tree bark?. Retrieved 13 February 2022, from <https://pursuit.unimelb.edu.au/articles/what-is-tree-bark>
- Merriam-Webster. (n.d.). Phlobatannin. In *Merriam-Webster.com dictionary*. Retrieved July 1, 2022, from <https://www.merriam-webster.com/dictionary/phlobatannin>
- Mondal, S. (2017). *Screening of bioactive compounds from Cucurbitaceae family edible plants of Bangladesh—Cucurbita pepo Linn.: A case study*. Anchor Academic Publishing.
- Mueller .M, & Tainter C. R. (2022). In: *StatPearls [Internet]: Escherichia Coli*. Treasure Island (FL):

StatPearls Publishing; 2022 Jan-. Available from:
<https://www.ncbi.nlm.nih.gov/books/NBK564298/>

- Mukherjee, P. K. (2010). *Infectious Diseases of the Eyes*. India: Lippincott Williams & Wilkins.
- Nawaz, H., Aslam, M., & Muntaha, S. (2019). Effect of Solvent Polarity and Extraction Method on Phytochemical Composition and Antioxidant Potential of Corn Silk. *Free Radicals and Antioxidants*, 9, 05-11. 10.5530/fra.2019.1.2.
- Nawaz, H., Shad, M., Rehman, N., Andaleeb, H. & Ullah, N. (2020). Effect of solvent polarity on extraction yield and antioxidant properties of phytochemicals from bean (*Phaseolus vulgaris*) seeds. *Brazilian Journal of Pharmaceutical Sciences*, 56. 10.1590/s2175-97902019000417129.
- NCI Thesaurus. (2022). Disk Diffusion Method. Retrieved 15 February 2022, from https://ncit.nci.nih.gov/ncitbrowser/ConceptReport.jsp?dictionary=NCI_Thesaurus&ns=ncit&code=C85595
- Omar, S., Lemonnier, B., Jones, N., Ficker, C., Smith, M. L., Neema, C., Towers, G. H. N., Goel, K., Arnason, J.T. (2000). Antimicrobial activity of extracts of eastern North American hardwood trees and relation to traditional medicine. *Journal of Ethnopharmacology*, 73(1-2), 161–170. doi:10.1016/s0378-8741(00)00294-4
- Ong, C. (2020). ANTIFUNGAL, ANTIBACTERIAL AND CYTOTOXIC ACTIVITIES OF NON-INDIGENOUS MEDICINAL PLANTS NATURALISED IN MALAYSIA. *Farmacia*, 68(4), 687-696. doi: 10.31925/farmacia.2020.4.14
- Othman, L., Sleiman, A., & Abdel-Massih, R. (2019). Antimicrobial Activity of Polyphenols and Alkaloids in Middle Eastern Plants. *Frontiers In Microbiology*, 10. doi: 10.3389/fmicb.2019.00911
- Panche, A. N., Diwan, A. D., & Chandra, S. R. (2016). Flavonoids: an overview. *Journal of nutritional science*, 5, e47. <https://doi.org/10.1017/jns.2016.41>
- Pásztor, Z., Mohácsiné, I., Gorbacheva, G., & Börcsök, Z. (2016). The Utilization of Tree Bark. *BioResources*, 11(3), 7859-7888.
- Patra A. K. (2012). An Overview of Antimicrobial Properties of Different Classes of Phytochemicals. *Dietary Phytochemicals and Microbes*, 1–32. https://doi.org/10.1007/978-94-007-3926-0_1
- Percival, S., & Williams, D. (2014). *Escherichia coli*. *Microbiology Of Waterborne Diseases*, 89-117. doi: 10.1016/b978-0-12-415846-7.00006-8
- Puupponen-Pimiä, R., Nohynek, L., Meier, C., Kähkönen, M., Heinonen, M., Hopia, A. & Oksman-Caldentey, K. (2001). Antimicrobial properties of phenolic compounds from berries. *Journal of Applied Microbiology*, 90, 494-507. <https://doi.org/10.1046/j.1365-2672.2001.01271.x>

- Radhia, A., Hanen, N., Abdelkarim, B. A., & Mohamed, N. (2018). Phytochemical screening, antioxidant and antimicrobial activities of *Erodium glaucophyllum* (L.) L'Hérit. *Journal of Biomedical Sciences*, 7(4).
- Raitanen, J.-E.; Järvenpää, E.; Korpinen, R.; Mäkinen, S.; Hellström, J.; Kilpeläinen, P.; Liimatainen, J.; Ora, A.; Tupasela, T.; & Jyske, T. (2020). Tannins of Conifer Bark as Nordic Piquancy—Sustainable Preservative and Aroma? *Molecules*, 25(3), 567. <https://doi.org/10.3390/molecules25030567>
- Sainath, R., Prathiba, J., & Malathi, R. (2009). Antimicrobial properties of the stem bark of *Saraca indica* (Caesalpinaceae). *European Review For Medical And Pharmacological Sciences*, (13), 371-374.
- Salem, M., Elansary, H., Elkelish, A., Zeidler, A., Ali, H., EL-Hefny, M., & Yessoufou, K. (2016). In vitro Bioactivity and Antimicrobial Activity of *Picea abies* and *Larix decidua* Wood and Bark Extracts. *BioResources*, 11(4), 9421-9437. Retrieved from <https://ojs.cnr.ncsu.edu/index.php/BioRes/article/view/10186>
- Shantabi, L., & Jagetia, G. C. (2015). Phytochemical profiling of Kam-sabut, *Croton caudatus* Geiseler. *Res Rev J Bot Sci*, 4, 5-14.
- Shi, J., Arunasalam, K., Yeung, D., Kakuda, Y., Mittal, G., & Jiang, Y. (2004). Saponins from edible legumes: chemistry, processing, and health benefits. *Journal of medicinal food*, 7(1), 67–78. <https://doi.org/10.1089/109662004322984734>
- Singh, R., Singh, S., Maharia, R., & Garg, A. (2015). Identification of new phytoconstituents and antimicrobial activity in stem bark of *Mangifera indica* (L.). *Journal Of Pharmaceutical And Biomedical Analysis*, 105, 150-155. doi: 10.1016/j.jpba.2014.12.010
- Sivagnanasundaram, P., & Karunanayake, K. (2015). Phytochemical Screening and Antimicrobial Activity of *Artocarpus heterophyllus* and *Artocarpus altilis* Leaf and Stem Bark Extracts. *OUSL Journal*, 9(0), 1-17. doi: 10.4038/ouslj.v9i0.7324
- Steenkamp, J. A., Steynberg, J. P., Brandt, E. V., Ferreira, D. , & Roux, D. G. (1985). Phlobatannins, a novel class of ring-isomerized condensed tannins. *Journal of the Chemical Society, Chemical Communications*, (23), 1678–. doi:10.1039/c39850001678
- Steiger, P., Börner, A., & Schweingruber, F. H. (2019). *Bark Anatomy of Trees and Shrubs in the Temperate Northern Hemisphere*. Germany: Springer International Publishing.
- Stokland, J. N., Siitonen, J., & Jonsson, B. G. (2012). *Biodiversity in Dead Wood*. United Kingdom: Cambridge University Press.
- Sugita, P., Amilia, R., Arifin, B., Rahayu, D., & Dianhar, H. (2020). TheE Phytochemical Screening Hexane and Methanol Extract of Sinyo Nakal (*Durata Repens*). *Asian Journal of Pharmaceutical and Clinical Research*. 196-200. 10.22159/ajpcr.2020.v13i8.38165.

- Taylor, T. A., & Unakal, C. G., (2022). In: *StatPearls [Internet]: Staphylococcus Aureus*. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK441868/>
- Tendencia, E. A. (2004). *Disk diffusion method - Laboratory Manual of Standardized Methods for Antimicrobial Sensitivity Tests for Bacteria Isolated from Aquatic Animals and Environment* (pp. 13-29). Tigbauan, Iloilo, Philippines: Aquaculture Department, Southeast Asian Fisheries Development Center.
- Tong, S. Y., Davis, J. S., Eichenberger, E., Holland, T. L., & Fowler, V. G., Jr (2015). Staphylococcus aureus infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clinical microbiology reviews*, 28(3), 603–661. <https://doi.org/10.1128/CMR.00134-14>
- Truong, D., Nguyen, D., Ta, N., Bui, A., Do, T., & Nguyen, H. (2019). Evaluation of the Use of Different Solvents for Phytochemical Constituents, Antioxidants, and *In Vitro* Anti-Inflammatory Activities of *Severinia buxifolia*. *Journal Of Food Quality*, 2019, 1-9. doi: 10.1155/2019/8178294
- USDA. Tannins. Retrieved 6 May 2022, from <https://www.fs.fed.us/wildflowers/ethnobotany/tannins.shtml#:~:text=Tannins%20are%20complex%20chemical%20substances,all%20parts%20of%20the%20globe.>
- Wadood, A., Ghufuran, M., Jamal, S. B., Naeem, M., Khan, A., & Ghaffar, R. (2013). Phytochemical analysis of medicinal plants occurring in local area of Mardan. *Biochem Anal Biochem*, 2(4), 1-4.
- Wagner, K., Roth, C., Willför, S., Musso, M., Petutschnigg, A., Oostingh, G., & Schnabel, T. (2019). Identification of Antimicrobial Compounds in Different Hydrophilic Larch Bark Extracts. *BioResources*, 14(3), 5807-5815.
- Wijaya, T. H., & Ekowati, N. (2021). The Antibacterial Activity of *Acanthus ilicifolius* L. n-Hexane Fraction. *Journal of Science and Technology Research for Pharmacy*, 1(2), 48-56.
- Xie, Y., Yang, W., Tang, F., Chen, X., & Ren, L. (2015). Antibacterial activities of flavonoids: structure-activity relationship and mechanism. *Current medicinal chemistry*, 22(1), 132–149. <https://doi.org/10.2174/0929867321666140916113443>
- Yang, Y., Laval, S., & Yu, B. (2021). Chemical Synthesis of Saponins. *Advances In Carbohydrate Chemistry And Biochemistry*, 63-150. doi: 10.1016/bs.accb.2021.10.001
- Yosef Sham, A., & Almasri, A. (2018). Bacteriostatic and Bactericidal Activity of Deer Musk on Multidrug Resistance Bacteria. *Pakistan Journal Of Biological Sciences*, 21(7), 331-339. doi: 10.3923/pjbs.2018.331.339
- Yusuf, A., Abdullahi, M., Aleku, G., Ibrahim, I., Alebiosu, C., & Yahaya, M. et al. (2018). Antimicrobial activity of stigmasterol from the stem bark of *Neocarya macrophylla*. *Journal Of Medicinal Plants For Economic Development*, 2(1), 1-5..

- Zhang, Q. W., Lin, L. G., & Ye, W. C. (2018). Techniques for extraction and isolation of natural products: a comprehensive review. *Chinese medicine*, 13, 20. <https://doi.org/10.1186/s13020-018-0177-x>
- Zhang, K., & Wong, J. (2011). Solvent-Based Extraction Techniques for the Determination of Pesticides in Food. *Comprehensive Sampling And Sample Preparation*, 245-261. doi: 10.1016/b978-0-12-381373-2.00136-8

APPENDICES

Appendix 1. Table replication of KB and KK extraction

Sample	1 st Extraction	2 nd Extraction	3 rd Extraction	Average
KBM	1.00%	0.93%	1.06%	1.00%
KBA	0.26%	1.02%	0.63%	0.64%
KBC	2.84%	1.50%	1.14%	1.83%
KBE	0.11%	0.76%	1.21%	0.70%
KBH	0.61%	0.52%	1.56%	0.89%
Total Average				1.01%
KKM	8.77%	8.25%	17.29%	11.44%
KKA	4.45%	8.40%	13.70%	8.85%
KKC	7.10%	4.49%	3.31%	4.97%
KKE	2.97%	4.19%	4.78%	3.98%
KKH	0.91%	0.99%	0.69%	0.86%
Total Average				6.02%

KB: *Kayu Bawang*, KK: *Kayu Kamlowelen*; The third letter represents the solvent by which the sample was extracted. M: Methanol, A: Acetone, C: Chloroform, E: Ethyl acetate, H: Hexane.


Appendix 2. Kayu Bawang Gisikisik Documentation



Stems and leaves of *Kayu Bawang*



Fine powder of *Kayu Bawang*



Maceration of *Kayu Bawang*



Extract of *Kayu Bawang*

Appendix 3. Kayu Teh Kamlowelen Documentation



Stems and leaves of *Kayu Kamlowelen*



Fine powder of *Kayu Kamlowelen*



Maceration of *Kayu Kamlowelen*



Extract of *Kayu Kamlowelen*