

References

- Bernard, F.-X., Morel, F., Camus, M., Pedretti, N., Barrault, C., Garnier, J., & Lecron, J.-C. (2012). Keratinocytes under Fire of Proinflammatory Cytokines: Bona Fide Innate Immune Cells Involved in the Physiopathology of Chronic Atopic Dermatitis and Psoriasis. *Journal of Allergy*, 2012, 1–10. <https://doi.org/10.1155/2012/718725>
- Bhalla, T. N., Saxena, R. C., Nigam, S. K., Misra, G., & Bhargava, K. P. (1980). Calophyllolide--a new non-steroidal anti-inflammatory agent. *The Indian journal of medical research*, 72, 762–765.
- Boer, M., Duchnik, E., Maleszka, R., & Marchlewicz, M. (2016). Structural and biophysical characteristics of human skin in maintaining proper epidermal barrier function. *Advances In Dermatology And Allergology*, 1, 1-5. <https://doi.org/10.5114/pdia.2015.48037>
- Cambie, R. C., & Ash, J. (1994). Fijian medicinal plants. CSIRO publishing.
- Cao, E., Chen, Y., Cui, Z., & Foster, P. R. (2003b). Effect of freezing and thawing rates on denaturation of proteins in aqueous solutions. *Biotechnology and Bioengineering*, 82(6), 684–690. <https://doi.org/10.1002/bit.10612>
- Cassien, M., Mercier, A., Thétiot-Laurent, S., Culcasi, M., Ricquebourg, E., Asteian, A., Herbette, G., Bianchini, J.-P., Raharivelomanana, P., & Pietri, S. (2021). Improving the antioxidant properties of *Calophyllum inophyllum* seed oil from French Polynesia: Development and biological applications of resinous ethanol-soluble extracts. *Antioxidants*, 10(2), 199. <https://doi.org/10.3390/antiox10020199>
- Catherine, J., & Roufosse, F. (2021). What does elevated TARC/CCL17 expression tell us about eosinophilic disorders? *Seminars in Immunopathology*, 43. <https://doi.org/10.1007/s00281-021-00857-w>
- Cayrol, C., & Girard, J.-P. (2017). Interleukin-33 (IL-33): A nuclear cytokine from the IL-1 family. *Immunological Reviews*, 281(1), 154–168. <https://doi.org/10.1111/imr.12619>
- Chamlin, S. L., Kao, J., Frieden, I. J., Sheu, M. Y., Fowler, A. J., Fluhr, J. W., Williams, M. L., & Elias, P. M. (2002). Ceramide-dominant barrier repair lipids alleviate childhood atopic dermatitis: Changes

in barrier function provide a sensitive indicator of disease activity. *Journal of the American Academy of Dermatology*, 47(2), 198–208. <https://doi.org/10.1067/mjd.2002.124617>

Cianferoni, A., & Spergel, J. (2014). The importance of TSLP in allergic disease and its role as a potential therapeutic target. *Expert Review of Clinical Immunology*, 10(11), 1463–1474. <https://doi.org/10.1586/1744666x.2014.967684>

Colombo, I., Sangiovanni, E., Maggio, R., Mattozzi, C., Zava, S., Corbett, Y., Fumagalli, M., Carlino, C., Corsetto, P. A., Scaccabarozzi, D., Calvieri, S., Gismondi, A., Taramelli, D., & Dell'Agli, M. (2017). HaCaT cells as a reliable in vitro differentiation model to dissect the inflammatory/repair response of human keratinocytes. *Mediators of Inflammation*, 2017, 1–12. <https://doi.org/10.1155/2017/7435621>

Coondoo, A., Phiske, M., Verma, S., & Lahiri, K. (2014). Side-effects of topical steroids: A long overdue revisit. *Indian Dermatology Online Journal*, 5(4), 416. <https://doi.org/10.4103/2229-5178.142483>

David Boothe, W., Tarbox, J. A., & Tarbox, M. B. (2017). Atopic dermatitis: Pathophysiology. *Advances in Experimental Medicine and Biology*, 21–37. https://doi.org/10.1007/978-3-319-64804-0_3

De Benedetto, A., Rafaels, N. M., McGirt, L. Y., Ivanov, A. I., Georas, S. N., Cheadle, C., Berger, A. E., Zhang, K., Vidyasagar, S., Yoshida, T., Boguniewicz, M., Hata, T., Schneider, L. C., Hanifin, J. M., Gallo, R. L., Novak, N., Weidinger, S., Beaty, T. H., Leung, D. Y. M., ... Beck, L. A. (2011). Tight Junction defects in patients with atopic dermatitis. *Journal of Allergy and Clinical Immunology*, 127(3). <https://doi.org/10.1016/j.jaci.2010.10.018>

Dębińska, A. (2021). New Treatments for Atopic Dermatitis Targeting Skin Barrier Repair via the Regulation of FLG Expression. *Journal of Clinical Medicine*, 10(11), 2506. <https://doi.org/10.3390/jcm10112506>

Deleuran, M., Hvid, M., Kemp, K., Christensen, G. B., Deleuran, B., & Vestergaard, C. (2012). IL-25 induces both inflammation and skin barrier dysfunction in atopic dermatitis. *New Trends in Allergy and Atopic Eczema*, 45–49. <https://doi.org/10.1159/000331871>

- Deng, C., Peng, N., Tang, Y., Yu, N., Wang, C., Cai, X., Zhang, L., Hu, D., Ciccia, F., & Lu, L. (2021). Roles of il-25 in type 2 inflammation and autoimmune pathogenesis. *Frontiers in Immunology*, 12. <https://doi.org/10.3389/fimmu.2021.691559>
- Dong, L., Cao, T. Q., Liu, Z., Tuan, N. Q., Kim, Y.-C., Sohn, J. H., Yim, J. H., Lee, D.-S., & Oh, H. (2022). Anti-inflammatory effects exerted by 14-methoxyalternate C from Antarctic fungal strain Pleosporales SP.. SF-7343 via the regulation of NF-KB and JAK2/STAT3 in HaCaT human keratinocytes. *International Journal of Molecular Sciences*, 23(23), 14642. <https://doi.org/10.3390/ijms232314642>
- Ebina-Shibuya, R., & Leonard, W. J. (2022). Role of thymic stromal lymphopoietin in allergy and beyond. *Nature Reviews Immunology*. <https://doi.org/10.1038/s41577-022-00735-y>
- Eisenberg, E., & Levanon, E. Y. (2013). Human housekeeping genes, revisited. *Trends in Genetics*, 29(10), 569–574. <https://doi.org/10.1016/j.tig.2013.05.010>
- Friday, J. B., & Okano, D. (2006). *Calophyllum inophyllum (kamani)*. Species profiles for Pacific Island agroforestry, 2(1), 1-17.
- Gallagher, S.R. and Wiley, E.A. (2012) '10', in Current Protocols Essential Laboratory Techniques. Hoboken, N.J: Wiley-Blackwell, p. 10.2.26-10.2.27.
- Gimalova, G. F., Karunas, A. S., Fedorova, Y. Y., Gumennaya, E. R., Levashova, S. V., Khismatullina, Z. R., Prans, E., Kōks, S., Etkina, E. I., & Khusnutdinova, E. K. (2014). Association of polymorphisms in the toll-like receptor genes with atopic dermatitis in the Republic of Bashkortostan. *Molecular Biology*, 48(2), 227–237. <https://doi.org/10.1134/s002689331402006x>
- Ginigini, J., Lecellier, G. J., Nicolas, M., Nour, M., Hnawia, E., Lebouvier, N., Herbette, G., Lockhart, P., & Raharivelomanana, P. (2019). Chemodiversity of *calophyllum inophyllum*l. oil bioactive components related to their specific geographical distribution in the South Pacific region. *PeerJ*, 7. <https://doi.org/10.7717/peerj.6896>

Ginwala, R. et al. (2019) 'Potential role of flavonoids in treating chronic inflammatory diseases with a special focus on the anti-inflammatory activity of Apigenin', *Antioxidants*, 8(2), p. 35.
<https://doi.org/10.3390/antiox8020035>.

Gruber, R., Elias, P. M., Crumrine, D., Lin, T.-K., Brandner, J. M., Hachem, J.-P., Presland, R. B., Fleckman, P., Janecke, A. R., Sandilands, A., McLean, W. H. I., Fritsch, P. O., Mildner, M., Tschachler, E., & Schmuth, M. (2011). Filaggrin genotype in ichthyosis vulgaris predicts abnormalities in epidermal structure and function. *The American Journal of Pathology*, 178(5), 2252–2263. <https://doi.org/10.1016/j.ajpath.2011.01.053>

Guttman-Yassky, E., Suárez-Fariñas, M., Chiricozzi, A., Nogales, K. E., Shemer, A., Fuentes-Duculan, J., Cardinale, I., Lin, P., Bergman, R., Bowcock, A. M., & Krueger, J. G. (2009). Broad defects in epidermal cornification in atopic dermatitis identified through genomic analysis. *Journal of Allergy and Clinical Immunology*, 124(6). <https://doi.org/10.1016/j.jaci.2009.09.031>

Ha, Y., Lee, W.-H., Jeong, J. W., Park, M., Ko, J.-Y., Kwon, O. W., Lee, J., & Kim, Y.-J. (2020). Pyropia yezoensis extract suppresses IFN-gamma- and TNF-alpha-induced proinflammatory chemokine production in HaCaT cells via the down-regulation of NF-KB. *Nutrients*, 12(5), 1238. <https://doi.org/10.3390/nu12051238>

Hadi, H. A., Tarmizi, A. I., Khalid, K. A., Gajdács, M., Aslam, A., & Jamshed, S. (2021). The epidemiology and global burden of atopic dermatitis: A narrative review. *Life*, 11(9), 936. <https://doi.org/10.3390/life11090936>

Hashimoto, S., Nakamura, K., Oyama, N., Kaneko, F., Tsunemi, Y., Saeki, H., & Tamaki, K. (2006). Macrophage-derived chemokine (MDC)/CCL22 produced by monocyte derived dendritic cells reflects the disease activity in patients with atopic dermatitis. *Journal of Dermatological Science*, 44(2), 93–99. <https://doi.org/10.1016/j.jdermsci.2006.08.004>

He, R., & Geha, R. S. (2010). Thymic stromal lymphopoitin. *Annals of the New York Academy of Sciences*, 1183(1), 13–24. <https://doi.org/10.1111/j.1749-6632.2009.05128.x>

- Homey, B., Alenius, H., Müller, A., Soto, H., Bowman, E. P., Yuan, W., McEvoy, L., Lauerma, A. I., Assmann, T., Büinemann, E., Lehto, M., Wolff, H., Yen, D., Marxhausen, H., To, W., Sedgwick, J., Ruzicka, T., Lehmann, P., & Zlotnik, A. (2002). CCL27–CCR10 interactions regulate T cell–mediated skin inflammation. *Nature Medicine*, 8(2), 157–165.
<https://doi.org/10.1038/nm0202-157>
- Howell, M. D., Kim, B. E., Gao, P., Grant, A. V., Boguniewicz, M., DeBenedetto, A., Schneider, L., Beck, L. A., Barnes, K. C., & Leung, D. Y. M. (2007). Cytokine modulation of atopic dermatitis filaggrin skin expression. *Journal of Allergy and Clinical Immunology*, 120(1), 150–155.
<https://doi.org/10.1016/j.jaci.2007.04.031>
- Idriss, H. T., & Naismith, J. H. (2000). TNF? and the TNF receptor superfamily: Structure-function relationship(s). *Microscopy Research and Technique*, 50(3), 184–195.
[https://doi.org/10.1002/1097-0029\(20000801\)50:3<184::aid-jemt2>3.0.co;2-h](https://doi.org/10.1002/1097-0029(20000801)50:3<184::aid-jemt2>3.0.co;2-h)
- Imai, Y., Yasuda, K., Sakaguchi, Y., Haneda, T., Mizutani, H., Yoshimoto, T., Nakanishi, K., & Yamanishi, K. (2013). Skin-specific expression of IL-33 activates group 2 innate lymphoid cells and elicits atopic dermatitis-like inflammation in mice. *Proceedings of the National Academy of Sciences*, 110(34), 13921–13926. <https://doi.org/10.1073/pnas.1307321110>
- Jain, K., Salamat-Miller, N., & Taylor, K. (2021). Freeze–thaw characterization process to minimize aggregation and enable drug product manufacturing of protein based therapeutics. *Scientific Reports*, 11(1). <https://doi.org/10.1038/s41598-021-90772-9>
- Kakinuma, T., Saeki, H., Tsunemi, Y., Fujita, H., Asano, N., Mitsui, H., Tada, Y., Wakugawa, M., Watanabe, T., Torii, H., Komine, M., Asahina, A., Nakamura, K., & Tamaki, K. (2003). Increased serum cutaneous T cell-attracting chemokine (CCL27) levels in patients with atopic dermatitis and psoriasis vulgaris. *Journal of Allergy and Clinical Immunology*, 111(3), 592–597.
<https://doi.org/10.1067/mai.2003.114>

Kantor, R., & Silverberg, J. I. (2016). Environmental risk factors and their role in the management of atopic dermatitis. *Expert Review of Clinical Immunology*, 13(1), 15–26.
<https://doi.org/10.1080/1744666x.2016.1212660>

Kapur, S., Watson, W., & Carr, S. (2018). Atopic dermatitis. *Allergy, Asthma & Clinical Immunology*, 14(S2). <https://doi.org/10.1186/s13223-018-0281-6>

Kezic, S., O'Regan, G. M., Yau, N., Sandilands, A., Chen, H., Campbell, L. E., Kroboth, K., Watson, R., Rowland, M., Irwin McLean, W. H., & Irvine, A. D. (2011). Levels of filaggrin degradation products are influenced by both Filaggrin genotype and atopic dermatitis severity. *Allergy*, 66(7), 934–940. <https://doi.org/10.1111/j.1398-9995.2010.02540.x>

Kim, B. E., Leung, D. Y. M., Boguniewicz, M., & Howell, M. D. (2008). Loricrin and involucrin expression is down-regulated by th2 cytokines through STAT-6. *Clinical Immunology*, 126(3), 332–337.
<https://doi.org/10.1016/j.clim.2007.11.006>

Kim, H. J., Baek, J., Lee, J. R., Roh, J. Y., & Jung, Y. J. (2018). Optimization of cytokine milieu to reproduce atopic dermatitis-related gene expression in HaCaT keratinocyte cell line. *Immune Network*, 18(2). <https://doi.org/10.4110/in.2018.18.e9>

Kim, J., Kim, B. E., & Leung, D. Y. (2019). Pathophysiology of atopic dermatitis: Clinical implications. *Allergy and Asthma Proceedings*, 40(2), 84–92. <https://doi.org/10.2500/aap.2019.40.4202>

Kim, K., Kim, H., & Sung, G. Y. (2022). An interleukin-4 and interleukin-13 induced atopic dermatitis human skin equivalent model by a skin-on-a-chip. *International Journal of Molecular Sciences*, 23(4), 2116. <https://doi.org/10.3390/ijms23042116>

Klonowska, J., Gleń, J., Nowicki, R., & Trzeciak, M. (2018). New cytokines in the pathogenesis of atopic dermatitis—new therapeutic targets. *International Journal of Molecular Sciences*, 19(10), 3086.
<https://doi.org/10.3390/ijms19103086>

Kolb, L., & Ferrer-Bruker, S. J. (2022). Atopic Dermatitis. Nih.gov; StatPearls Publishing.
<https://www.ncbi.nlm.nih.gov/books/NBK448071/>

- Kuo, I.-H., Carpenter-Mendini, A., Yoshida, T., McGirt, L. Y., Ivanov, A. I., Barnes, K. C., Gallo, R. L., Borkowski, A. W., Yamasaki, K., Leung, D. Y., Georas, S. N., De Benedetto, A., & Beck, L. A. (2013). Activation of epidermal toll-like receptor 2 enhances tight junction function: Implications for atopic dermatitis and skin barrier repair. *Journal of Investigative Dermatology*, 133(4), 988–998. <https://doi.org/10.1038/jid.2012.437>
- Leung, D., & Guttman-Yassky, E. (2014). Deciphering the complexities of atopic dermatitis: Shifting paradigms in treatment approaches. *Journal Of Allergy And Clinical Immunology*, 134(4), 769-779. <https://doi.org/10.1016/j.jaci.2014.08.008>
- Margolis, D., Apter, A., Gupta, J., Hoffstad, O., Papadopoulos, M., & Campbell, L. et al. (2012). The persistence of atopic dermatitis and filaggrin (FLG) mutations in a US longitudinal cohort. *Journal Of Allergy And Clinical Immunology*, 130(4), 912-917. <https://doi.org/10.1016/j.jaci.2012.07.008>
- Matsunaga, M. C., & Yamauchi, P. S. (2016). IL-4 and IL-13 Inhibition in Atopic Dermatitis. *Journal of drugs in dermatology: JDD*, 15(8), 925-929.
- Merivaara, A., Zini, J., Koivunotko, E., Valkonen, S., Korhonen, O., Fernandes, F. M., & Yliperttula, M. (2021). Preservation of biomaterials and cells by freeze-drying: Change of paradigm. *Journal of Controlled Release*, 336, 480–498. <https://doi.org/10.1016/j.jconrel.2021.06.042>
- Miyano, K., & Tsunemi, Y. (2020). Current treatments for atopic dermatitis in Japan. *The Journal of Dermatology*, 48(2), 140–151. <https://doi.org/10.1111/1346-8138.15730>
- Nguyen, M.N. et al. (2021) ‘Purification trials of tamanu (*calophyllum inophyllum* L.) oil’, *OCCL*, 28, p. 53. <https://doi.org/10.1051/ocl/2021042>.
- Nutten, S. (2015). Atopic Dermatitis: Global Epidemiology and Risk Factors. *Annals of Nutrition and Metabolism*, 66(Suppl. 1), 8–16. <https://doi.org/10.1159/000370220>
- Nygaard, U., Bogaard, E., Niehues, H., Hvid, M., Deleuran, M., Johansen, C., & Vestergaard, C. (2017). The “alarmins” HMBG1 and IL-33 downregulate structural skin barrier proteins and impair

epidermal growth. Acta Dermato Venereologica, 97(3), 305–312.

<https://doi.org/10.2340/00015555-2552>

Nowicki, R., Trzeciak, M., Wilkowska, A., Sokołowska-Wojdyło, M., Ługowska-Umer, H., Barańska-Rybak, W., Kaczmarski, M., Kowalewski, C., Kruszewski, J., Maj, J., Silny, W., Śpiewak, R., & Petranyuk, A. (2015). Special paper atopic dermatitis: Current treatment guidelines. statement of the experts of the dermatological section, polish Society of Allergology, and the allergology section, polish Society of Dermatology. Advances in Dermatology and Allergology, 4, 239–249. <https://doi.org/10.5114/pdia.2015.53319>

Oh, J.-H., Kim, S.-H., Kwon, O.-K., Kim, J.-H., Oh, S.-R., Han, S.-B., Park, J.-W., & Ahn, K.-S. (2022). Purpurin suppresses atopic dermatitis via TNF- α /IFN- γ -induced inflammation in HaCaT cells. International Journal of Immunopathology and Pharmacology, 36, 3946320221111135. <https://doi.org/10.1177/03946320221111135>

Prabakaran, K., & Britto, S. J. (2012). Biology, agroforestry and medicinal value of *Calophyllum inophyllum* L.(clusiacea): A review. International Journal of Natural Products Research, 1(2), 24-33.

Pribowo, A., Girish, J., Gustiananda, M., Nandhira, R., & Hartrianti, P. (2021). Potential of Tamanu (*Calophyllum inophyllum*) Oil for Atopic Dermatitis Treatment. Evidence-Based Complementary And Alternative Medicine, 2021, 1-9. <https://doi.org/10.1155/2021/6332867>

Pyun, B. Y. (2015). Natural history and risk factors of atopic dermatitis in children. Allergy, Asthma & Immunology Research, 7(2), 101. <https://doi.org/10.4168/aaair.2015.7.2.101>

Raharivelomanana, P., Ansel, J.-L., Lupo, E., Mijouin, L., Guillot, S., Butaud, J.-F., Ho, R., Lecellier, G., & Pichon, C. (2018). Tamanu oil and skin active properties: From traditional to modern cosmetic uses. OCL, 25(5). <https://doi.org/10.1051/ocl/2018048>

Rahman, Md.M. et al. (2021) 'Role of phenolic compounds in human disease: Current knowledge and future prospects', Molecules, 27(1), p. 233. <https://doi.org/10.3390/molecules27010233>.

- Saeki, H., & Tamaki, K. (2008). Role of TARC/CCL17 and CTACK/CCL27 in the pathogenesis of atopic dermatitis. *Current Immunology Reviews*, 4(4), 221–229.
<https://doi.org/10.2174/157339508786447878>
- Sandilands, A., Sutherland, C., Irvine, A. D., & McLean, W. H. (2009). Filaggrin in the frontline: Role in skin barrier function and disease. *Journal of Cell Science*, 122(9), 1285–1294.
<https://doi.org/10.1242/jcs.033969>
- Sano, Y., Masuda, K., Tamagawa-Mineoka, R., Matsunaka, H., Murakami, Y., Yamashita, R., Morita, E., & Katoh, N. (2013). Thymic stromal lymphopoietin expression is increased in the horny layer of patients with atopic dermatitis. *Clinical and Experimental Immunology*, 171(3), 330–337.
<https://doi.org/10.1111/cei.12021>
- Saravanan, R., Dhachinamoorthi, D., Senthilkumar, K., & Thamizhvanan, K. (2011). Antimicrobial activity of various extracts from various parts of *Calophyllum inophyllum* L.
- Schmitt, J., Schmitt, N., & Meurer, M. (2007). Cyclosporin in the treatment of patients with atopic eczema ? A systematic review and meta-analysis. *Journal of the European Academy of Dermatology and Venereology*. <https://doi.org/10.1111/j.1468-3083.2006.02023.x>
- Shimada, Y., Takehara, K., & Sato, S. (2004). Both th2 and th1 chemokines (TARC/CCL17, MDC/CCL22, and MiG/CXCL9) are elevated in sera from patients with atopic dermatitis. *Journal of Dermatological Science*, 34(3), 201–208. <https://doi.org/10.1016/j.jdermsci.2004.01.001>
- Silverberg, N. B., & Silverberg, J. I. (2015). Inside out or outside in: does atopic dermatitis disrupt barrier function or does disruption of barrier function trigger atopic dermatitis? *Cutis*, 96(6), 359-361. <https://pubmed.ncbi.nlm.nih.gov/26761930/>
- Skrypina, N. (2003). Total RNA suitable for molecular biology analysis. *Journal of Biotechnology*, 105(1–2), 1–9. [https://doi.org/10.1016/s0168-1656\(03\)00140-8](https://doi.org/10.1016/s0168-1656(03)00140-8)
- Sroka-Tomaszewska, J., & Trzeciak, M. (2021). Molecular mechanisms of atopic dermatitis pathogenesis. *International Journal of Molecular Sciences*, 22(8), 4130.
<https://doi.org/10.3390/ijms22084130>

Tamagawa-Mineoka, R., & Katoh, N. (2020). Atopic dermatitis: Identification and management of complicating factors. International Journal of Molecular Sciences, 21(8), 2671. <https://doi.org/10.3390/ijms21082671>

Tanoto, K. (2022). Identification and Flavonoids in Ethanol-Soluble Calophyllum Inophyllum Oil from Indonesia [Unpublished manuscript]. Faculty of Pharmacy, Indonesia International Institute for Life Sciences.<http://repository.i3l.ac.id/jspui/handle/123456789/740>

Tau, G., & Rothman, P. (1999). Biologic functions of the IFN-gamma receptors. Allergy, 54(12), 1233–1251. <https://doi.org/10.1034/j.1398-9995.1999.00099.x>

Thomsen, S. F. (2014). Atopic dermatitis: Natural history, diagnosis, and treatment. ISRN Allergy, 2014, 1–7. <https://doi.org/10.1155/2014/354250>

Urban, K., Chu, S., Giese, R. L., Mehrmal, S., Uppal, P., Nedley, N., & Delost, G. R. (2021). The global, regional, and national burden of atopic dermatitis in 195 countries and territories: An Ecological Study from the global burden of disease study 2017. JAAD International, 2, 12–18. <https://doi.org/10.1016/j.jdin.2020.10.002>

Vestergaard, C., Deleuran, M., Gesser, B., & Larsen, C. G. (2004). Thymus- and activation-regulated chemokine (TARC/CCL17) induces a th2-dominated inflammatory reaction on intradermal injection in mice. Experimental Dermatology, 13(4), 265–271. <https://doi.org/10.1111/j.0906-6705.2004.00149.x>

Walterscheid, J. P., Nghiem, D. X., Kazimi, N., Nutt, L. K., McConkey, D. J., Norval, M., & Ullrich, S. E. (2006). cis -urocanic acid, a sunlight-induced immunosuppressive factor, activates immune suppression via the 5-HT 2a receptor. Proceedings of the National Academy of Sciences, 103(46), 17420–17425. <https://doi.org/10.1073/pnas.0603119103>

Wirnsberger, G., Hebenstreit, D., Posselt, G., Horejs-Hoeck, J., & Duschl, A. (2006). IL-4 induces expression of TARC/CCL17 VIA two STAT6 binding sites. European Journal of Immunology, 36(7), 1882–1891. <https://doi.org/10.1002/eji.200635972>

Wang, R. et al. (2022) 'Biologically synthesized rosa rugosa-based gold nanoparticles suppress skin inflammatory responses via MAPK and NF-KB signaling pathway in TNF- α /IFN- γ -induced HaCaT keratinocytes', ACS Omega, 7(40), pp. 35951–35960.
<https://doi.org/10.1021/acsomega.2c04832>.

Yang, G., Seok, J., Kang, H., Cho, Y., Lee, H., & Lee, J. (2020). Skin Barrier Abnormalities and Immune Dysfunction in Atopic Dermatitis. International Journal Of Molecular Sciences, 21(8), 2867.
<https://doi.org/10.3390/ijms21082867>

Yimdjio, M. C., Azebaze, A. G., Nkengfack, A. E., Meyer, A. M., Bodo, B., & Fomum, Z. T. (2004). Antimicrobial and cytotoxic agents from Calophyllum inophyllum. Phytochemistry, 65(20), 2789–2795. <https://doi.org/10.1016/j.phytochem.2004.08.024>